PHYSICAL SCI.LIB.

TC 824 C2 A2 No.202-78



lifornia roes Agency

Lof

UNIVERSITY OF CALIFORNIA DAVIS

MAR 2 8 1980

GOV'T. DOCS. - LIBRARY



Conditions and Events in ornia

ar 1977-78

2-78 1979



COVER PHOTOGRAPH: Water cascading in the upper Sacramento River, about mid December 1977, provided the first sign of relief from two consecutive record dry water years of 1976 and 1977.

Department of Water Resources

Bulletin 202-78

Water Conditions and Flood Events in California

Water Year 1977-78

December 1979

Huey D. Johnson Secretary for Resources

The Resources Agency

Edmund G. Brown Jr.

State of California

Ronald B. Robie Director

Department of Water Resources

FIGURE 1. COUNTIES PROCLAIMED DISASTER AREAS DURING WATER YEAR 1977-78



FOREWORD

Water year 1977-78 (October 1, 1977 through September 30, 1978) marked the return of normal water supplies and storage in California. Ten weeks after the water year began, relatively heavy December storms brought long-awaited relief from the record drought conditions of the two previous years. The heavy precipitation began in mid December 1977, and continued almost without interruption until January 20, 1978. On February 1, 1978, the Department forecast above-average water year runoff if median conditions of precipitation prevailed for the remainder of the season. By the end of the season on May 1, 1978, the water year forecast of statewide snowmelt runoff was 170 percent of average.

Bulletin 202-78 combines general information on statewide water conditions and significant flooding events of the water year 1977-78. The Bulletin covers weather patterns of significant storm periods; and information on precipitation, snowpack, unimpaired runoff, and reservoir storage. The Bulletin also includes hydrographs of stream stages and reservoir operations, weir overflow graphs, tabulations of peak streamflow and stages, and other data on flood events during the year.

As part of the Department's continuing effort to reduce costs, Bulletin 202 will not be published again. The data contained in this Bulletin will be compiled, however, and available by request.

This report was prepared from information provided by the Department of Water Resources, National Weather Service, U. S. Geological Survey, U. S. Army Corps of Engineers, U. S. Water and Power Resources Service, and many other agencies, public and private, to whom we wish to express our thanks.

Ronald B. Robie, Director
Department of Water Resources
The Resources Agency

The Resources Agency State of California



CONTENTS

	Page
FOREWORD	iii
ORGANIZATION, DEPARTMENT OF WATER RESOURCES	vii
CONVERSION FACTORS - METRIC TO ENGLISH SYSTEM MEASUREMENT	viii
CHAPTER I. WEATHER	1
CHAPTER II. WATER SUPPLY	12 13 20 21 22
CHAPTER III. FLOOD EVENTS	25
APPENDIXES	
Appendix A: Sacramento River Crest and Weir Overflow Records (Figures)	43 44 45 46 47 48 49 50
Appendix B: Peak Flows and Stages at Selected Streams and Stations in California (Tables)	51 51
Peak Flows and Stages North Coastal Area San Francisco Bay Area Central Coastal Area South Coastal Area Central Valley Area Northern Lahontan Area Southern Lahontan Area Legend.	52 54 58 60 64 74 76

CONTENTS (Continued)

		Page
	FIGURES	
1	Counties Proclaimed Disaster Areas During Water	ii
2	Year 1977-78	ix
	October 1, 1977 - September 30, 1978	x
4	Water Year Precipitation in Percent of Normal, October 1, 1977 - September 30, 1978	6
5	October 1977 - April 1978	7
6 7	Water Content of Snowpack Accumulation in Percent	8
8	of April 1 Average	10 11
9 10 11	Unimpaired Runoff, 1977-78	13 16 22
12	Northern California Reference Map for Hydrographs, Figures 13-16	37
13 14	Hydrographs of Shasta Lake and Sacramento River Hydrographs of Lake Oroville and Folsom Lake	38 39
15 16	Hydrographs of Smith and Trinity Rivers	40 41
	TABLES	
1	Precipitation Amounts at Selected Stations During	5
2	Water Year 1977-78	14 18
3	Storage in Major Reservoirs	20
5	Comparisons of Water Supply Forecasts with Observed Unimpaired Runoff	21

Capies of this bulletin at \$3.00 each may be ordered fram: State of California
DEPARTMENT OF WATER RESOURCES
P. O. Box 388
Socramento, California 95802

Make checks payable to: DEPARTMENT OF WATER RESOURCES

California residents add 6% sales tax.

State of California EDMUND G. BROWN JR., Governor

The Resources Agency HUEY D. JOHNSON, Secretary for Resources

DEPARTMENT OF WATER RESOURCES RONALD B. ROBIE, Director

GERALD H. MERAL

MARY ANNE MARK

KEN YOSHIKAWA

ROBERT W. JAMES

Deputy Director	Deputy Director	Deputy Director
СН	ARLES R. SHOEMAKER Deputy Director	
DIVISI	ON OF FLOOD MANAGEMENT	
C. A. McCULLOUGH		Chief
This Bulletin v	was prepared under the di	rection
	of	
ALVIN J. BROWN DONALD H. NEUDECK	Chief, Sn	ow Surveys Branch Operations Branch
	by	
HOWARD J. SULLIVAN	Senior Meteorologis Water Resources Eng Mainten Water Resou	ineering Associate ance Supervisor II
	Assisted by	
William B. Chan George W. Patrick William D. Harrison . Jess C. Bringham	Water Resou Assistant Enginee Assistant Contr Water Resources Eng Water Resources S	r, Water Resources ol System Engineer ineering Associate rces Technician II
Draftin	ng under the direction of	

. . . Senior Delineator, Water Resources

CONVERSION FACTORS

Metric to Customary System of Measurement

Quantity	Metric Unit	Multiply by	To get customary equivalent
Length	millimetres (mm)	0.03937	inches (in)
	centimetres (cm) for snow depth	0.3937	inches (in)
	metres (m)	3.2808	feet (ft)
	kilometres (km)	0.62139	miles (m)
Area	square millimetres (mm²)	0.00155	square inches (in ²)
	square metres (m ²)	10.764	square feet (ft ²)
	hectares (ha)	2.4710	acres (ac)
	square kilometres (km²)	0.3861	square miles (mi ²)
Volume	litres (1)	0.26417	gallons (gal)
	megalitres	0.26417	million gallons (10 ⁶ gal)
	cubic metres (m ³)	35,315	cubic feet (ft ³)
	cubic metres (m ³)	1.308	cubic yards (yd ³)
	cubic metres (m ³)	0.0008107	acre-feet (ac-ft)
	cubic dekametres (dam ³)	0.8107	a cre-feet (ac-ft)
	cubic hectometres (hm ³)	0.8107	thousands of acre-feet
	cubic kilometres (km ³)	0.8107	millions of acre-feet
Flow	cubic metres per second (m ³ /s)	35.315	cubic feet per second (ft ³ /s)
	litres per minute (1/min)	0.26417	gallons per minute (gal/min)
	litres per day (I/day)	0.26417	gallons per day)(gal/day)
	megalitres per day (MI/day)	0.26417	million gallons per day (mgd)
	cubic metres per day (m ³ /day)	0.0008107	acre-feet per day
Mass	kilograms (kg)	2.2046	pounds (Ib)
	tonne (t)	1.1023	tons (short, 2,000 lb)
Velocity	metres per second (m/s)	3.2808	feet per second (ft/s)
Power	kilowatts (kW)	1.3405	horsepower (hp)
Pressure	kilopascals (kPa)	0.145054	pounds per square inch (psi)
	kilopascals (kPa)	0.33456	feet head of water
Specific capacity	litres per minute per metre drawdown	0.08052	gallons per minute per foot drawdown
Concentration	milligrams per litre (mg/l)	1.0	parts per million
Electrical conductivity	microsiemens per centimetre ($\mu S/cm$)	1.0	micromho per centimetre
Temperature	degrees Celsius (°C)	(1.8 × °C) + 32	degree Fahrenheit (F)

FIGURE 2. HYDROLOGIC BASINS OF CALIFORNIA





FIGURE 3. WATER YEAR PRECIPITATION IN INCHES (MILLIMETERS)

OCTOBER 1. 1, 1977 - SEPTEMBER 30, 1978



CHAPTER I - WEATHER

The weather patterns of the season 1977-78 marked a decisive change from the drought patterns of the two previous seasons. Once again weather fronts with thick cloud layers and ample moisture supply moved over California and dumped generous quantities of rain over the State. One professional weather watcher has called the winter 1977-78 for the West "the warm, drought-breaking winter of '78." The weather maps no longer showed a high-pressure ridge entrenched over the West that could divert storms away from California. Instead, storm systems moved across the Pacific on a wide path at mid latitudes to strike California broadside. Furthermore, numerous systems formed in the eastern Pacific at low latitudes and tracked northeastward toward California, particularly Southern California. All five months from December 1977 through April 1978 brought abovenormal precipitation to the State. The data in Table 1 illustrate how the fall-winter-spring seasons brought abundant precipitation at representative stations throughout the State.

In our discussion of the weather events of 1977-78, we will refer to two semipermanent atmospheric pressure systems, the so-called Aleutian low and the Pacific high.* The former is the low-(atmospheric) pressure center located in the north Pacific near the Aleutian Island chain (about Latitude $50^{\circ}N$), and the latter is the high-pressure center located in the eastern Pacific (Latitude 30° to 35°N). These pressure centers are important components of the air-circulation pattern over the Pacific sector of the northern hemisphere. In a broad sense, the Pacific high is strong (higher central pressure) and the Aleutian low relatively weak in the summer months, whereas in the winter months the opposite is true: the Aleutian low is strong and dominant (lower pressure) and the Pacific high is weak (pressure not as high as in summer). The pressure values on the daily weather maps averaged over a period of time (such as a month or season) define the strenght and location of these two pressure centers, and the weather phenomena (rain, clouds, sky condition, wind, etc.), in turn, are related to the defined patterns.

For convenience, the following discussion of the weather in 1977-78 has been broken down into monthly periods:

^{*} The meteorologist customarily draws lines of equal atmospheric pressure on the weather maps to delineate the low-and-high-pressure systems (or centers).

November. In November the westerly winds over the eastern Pacific began to increase, and the Aleutian low deepened below its normal November value. The Pacific Northwest and the northwestern corner of California began to experience weather fronts, and the resultant precipitation for the month was above normal for those areas mentioned. The storm tracks (i.e., the trajectories of the low-pressure centers) during November set up a favorable trend for storm events in the coming months to affect California.

<u>December</u>. December brought the further strengthening of the <u>Pacific</u> westerlies. Although the Aleutian low did continue below normal, the additional development took place whereby the Pacific high-pressure cell was weakened and was displaced to the southeast of its normal location. As a result, pressures west of California were at least 6-8 millibars* below normal; this permitted more frequent southerly winds to bring clouds and moisture over the State. Storm events in December were numerous, especially in the latter half of the month, and affected the entire State.

One noteworthy weather event (but not involving water) occurred on December 20. A very strong high-pressure cell developed suddenly during the night of December 19-20 over the Great Basin. This activated a very strong downslope wind through the Tehachapi canyons into the southern San Joaquin Valley. The wind gusts are estimated to have reached 160 kilometres per hour (100 m.p.h.). The strong, destructive winds mainly affected Kern County, and extensive damage to buildings and agriculture resulted from the blowing sand and dust. The hardest hit area was the Arvin-Lamont-Edison-Bakersfield sector. Large volumes of dust were carried by upper level southerly winds as far north as the Oregon-California border. A month later, snow surveyors on Mt. Shasta reported finding a layer of dust embedded in the snowpack that had been deposited in the December 20 storm.

Northern and Central California benefited from the December storms. Reservoirs began to accumulate runoff. For Southern California the storms came after December 16. The one important storm that helped boost the monthly totals significantly in Southern California occurred in the period December 23-29. This was a low-latitude storm which had developed in the eastern Pacific.

Overall, the monthly accumulations over the State were well above normal, with Southern California and the southern Sierra Nevada receiving amounts in excess of 200 percent of normal.

^{* 1} millibar = 0.03 inches mercury = 0.75 millimetres mercury = 0.10 kilopascal (SI unit of pressure).

January. In January 1978, the Aleutian low-pressure center was 10-12 millibars below normal and was located in its normal location (latitude 50° N, longitude 175° E). The eastern Pacific highpressure center was weakened and was displaced southward to 25° north latitude. An active storm track in the Pacific brought frequent frontal passages over California with accompanying precipitation. Precipitation totals over the State were well above normal -- many areas received 200 to 300 percent of normal, including the mountainous areas from which water supplies are primarily obtained. The plots of reservoir storage at major reservoirs, such as Shasta, Oroville, and Folsom, showed their most dramatic (sharp) rise in response to the repeated storm events. Snow levels in the January storms were in the intermediate elevation range -- 1 200 to 1 500 metres (4,000 to 5,000 feet), thus initiating the buildup of a deep snowpack in the mountains. For example, Lodgepole, in the Kaweah River drainage, started the month with 46 centimetres (18 inches) on the ground and had a depth of 206 centimetres (81 inches) by January 19.

February. February 1978 brought the continuation of a wet weather pattern for California. Both the Aleutian low-pressure and Pacific high-pressure centers were displaced southeastward from their normal locations. Frequent and intense migratory storms in the Pacific moved eastward along the southern periphery of the Aleutian low between 350 north latitude. Frontal systems moved through California and brought the accumulated monthly rainfall amounts to 1-1/2 to 3 times the normal over the State. Southern California was struck by a series of storms (extending into early March) that formed along the 30° latitude circle (about 1 100 kilometres [600 nautical miles] north of Hawaii) and tracked east-ward on a southerly path. The first storm affecting Southern California developed over the ocean on February 8 and continued into the 9th and 10th. Torrential rainfall in the Big Tujunga Canyon of the San Gabriel Mountains created a flood wave that brought destruction and loss of life to the community of Hidden Springs. Many stations in the mountains above the 600metre (1 968-foot) level had two-day totals exceeding 250 millimetres (10 inches). One of the highest two-day totals occurred at Crystal Lake in Los Angeles County where 432 millimetres (17 inches) was reported.

Snowpack accumulation in the mountains continued during the February storms. By the end of the month, the snow water storage had already exceeded the average total seasonal accumulation in all river basins south of the Pit River Basin. The water supply outlook for the summer of 1978 appeared good.

 $\frac{\mathrm{March}}{\mathrm{west}}$. In March 1978, the Aleutian low center was displaced southwest of its normal position, and pressures around the center were 8 to 10 millibars below normal. The Pacific high was displaced about 10° longitude west of its normal position. Frequent storm systems moved across the Pacific in the latitude band between 35° north and 45° north. The sharp fronts associated with the migratory low-pressure centers swept into California and released above-normal precipitation over the State (with the exception of the extreme North Coast). The rainy period covered generally the first part of the month through March 12, and March 21-23 and 30-31.

Southern California received a series of storms that began February 28 and continued into the first two days of March. These storms, like the one of February 9-10, had a southerly origin in the Pacific and tracked into Southern California. The two-day intensity of this storm almost equaled the one of February 9-10. Two stations reporting two-day amounts in the 305-330 millimetre range (12 to 13 inches) were Lytle Creek Ranger Station in San Bernardino County and Mt. Wilson 2 in Los Angeles County.

Another storm that followed closely had its heaviest concentration in the two-day period March 4-5 in the Southern California area. Lytle Creek Ranger Station reported a two-day total of 273 millimetres (10.76 inches). Property damage resulted from the torrential downpours and mudslides, and portions of the coast experienced heavy surf conditions, especially at Malibu.

TABLE 1. PRECIPITATION AMOUNTS AT SELECTED STATIONS DURING WATER YEAR 1977-78

HYDROLOGIC BASIN	FLEV	T 101					PR EC I	PITATION FOR	SELECTED	PER 1005				
HYDROLOGIC BASIN	ELEV,	ITON	0c1	Nov.	1977 Percent	MIIII-	inter Qua	rter	Spring Quarter			Water Year		
Station	Hetres	Feet	metres	Inches	Norma i	metres	Inches	Norma1	metres	Inches	Percent Normal	metres	Inches	Percent Normal
NORTH COAST Gasquet RS Eureka Fort Bragg	117 18 24	384 60 80	723 185 158	28.46 7.30 6.21	129 84 79	1 462 436 665	57.57 17.18 26.20	126 92 129	523 198 318	20.59 7.80 12.52	92 77 133	2 962 913 1 218	116.63 35.96 47.95	124 92 123
SACRAMENTO Shasta Dam Blue Canyon Sacramento El	328 1 610 26	1,075 5,280 84	169 193 58	6.66 7.61 2.27	60 64 75	1 425 1 257 407	56.09 44.50 16.03	180 151 157	671 703 165	26.42 27.68 6.50	181 169 148	2 400 2 313 639	94.47 91.05 25.17	159 144 140
SAN JOAQUIN Fresno AP	100	328	12	.47	29	269	10.59	182	180	7.10	220	488	19.21	178
TULARE LAKE Grant Grove Bakersfield	2 013 151	6,600 495	88	3.45	46 10	1 048	41.26	185 255	519 74	20.45	153 149	1 804	71.01 11.42	162 188
SAN FRANCISCO BAY Napa State Hospital Kentfield San Francisco CI	18 39 40	60 128 130	213 314 54	8.39 12.36 2.13	295 146 60	526 889 331	20.72 35.01 13.04	151 124 111	239 340 229	9.41 13.38 9.02	163 132 191	999 1 590 620	39.35 62.58 24.39	162 131 120
CENTRAL COAST Monterey San Luis Obispo Santa Barbara	117 96 2	385 315 5	17 8 0	.68 .33 0	22 11 0	442 888 595	17.41 34.96 23.43	175 262 218	272 316 344	10.69 12.46 13.53	229 231 313	741 1 245 968	29.19 49.00 38.12	159 222 215
SOUTH COASTAL AREA* Mt. Wilson 2 Los Angeles CC Sen Diego AP	1 741 78 4	5,709 257 13	18 2 14	.70 .08	15 4 34	1 291 541 261	50,81 21,31 10,26	266 241 181	652 249 147	25.65 9.79 5.77	323 290 239	2 029 802 439	79.88 31.57 17.30	245 217 174
NORTH LANONTAN Susanville AP	1 264	4,146	28	1,10	40	296	11.65	152	104	4.10	132	452	17.81	120
SOUTH LANONTAN Bishop AP Barstow	1 253 659	4,108 2,160	1 0.8	.05	6 4	217 144	8.54 5.67	271 346	47 69	1.86	172 342	280 226	11.01	194
COLORAGO RIVER Palm Springs Imperial	130 -20	425 -64	0 7	0 .29	0 71	178 65	7.02 2.54	224 227	52 12	2.05 .48	233 155	241 96	9.48 3.78	17F 147

^{*} South Coastal Area Includes the Los Angeles, Santa Ana, and San Blego Mydrologic Basins

Metric Equivalents: 1 inch = 25.4 millimetres (mm) 1 foot = 0.305 metre (m)

FIGURE 4. WATER YEAR PRECIPITATION IN PERCENT OF NORMAL OCTOBER 1, 1977 - SEPTEMBER 30, 1978






FIGURE 5. PRECIPITATION ACCUMULATION AT SELECTED STATIONS OCTOBER 1977 - APRIL 1978



IN







FIGURE 7. WATER CONTENT OF SNOWPACK ACCUMULATION IN PERCENT OF APRIL 1 AVERAGE



Intermittent heavy precipitation, which began in mid-December 1977 and continued through January 20, 1978, provided by February 1, 1978 a snowpack water content well above normal in all watersheds except the Pit River Basin. February 1 snow surveys indicated that snowpack water content of the Pit River Basin was 90 percent of the February 1 average; and the southern Sierra snowpack exceeded 150 percent of average. The February 1, 1978 Department forecast for the water year runoff throughout the Sierra varied from 115 to 140 percent of average, based on average precipitation and snowpack accumulation during the remainder of the season.


The March 1, 1978 snow surveys indicated that all river basins except the Pit River Basin had exceeded the average total seasonal accumulation, which normally occurs about April 1.

Snowpack measurements for March indicated that melting occurred at lower and mid elevations during March, but that by the month's end, all watersheds had a net increase in snowstored water. NASA satellite imagery for the southern Sierra watersheds showed an increase of 45 percent in snowcovered area for water year 1978 and over the record dry conditions of water year 1977.

The normal beginning of the snowmelt season in the Sierra Nevada is April 1. Low temperatures above 2 000 metres (6,500 feet) caused water accumulation to exceed snowmelt above this level during April in most areas. On May 1, 1978, a near-record pack existed, especially in the southern Sierra. Lower average temperatures until mid July were reflected by end-of-July satellite data on snowcover that indicated a substantial snowpack was left in protected high elevation portions of the southern Sierra watersheds

Seasonal snowpack accumulation curves are shown in Figure 7, and April 1 snowpack, in percent of average for individual river basins, is depicted in Figure 8.

FIGURE 8. SNOWPACK IN PERCENT OF AVERAGE APRIL 1, 1978



CHAPTER II - WATER SUPPLY

UNIMPAIRED RUNOFF

Unimpaired runoff during the 1977-78 water year for California was 158 percent of normal, compared to a record low of 24 percent of normal for 1976-1977. Relief from two consecutive years of record low runoff came in mid-December of 1977. Intermittent heavy rains during January 1978 resulted in runoff that was 200 percent of normal in the Central Valley, and which averaged almost 220 percent of normal statewide. Statewide, runoff continued to be normal and above normal for the remainder of the water year 1977-78.

Runoff from interior basins ranged from a high of 244 percent of average for the Kern River to 131 percent of average for the Feather and Yuba Rivers. San Francisco Bay hydrologic basin and Central Coast basin values were 167 and 295 percent of normal, respectively, compared to 1 and 5 percent of normal for the previous water year. Water year percentages for major hydrologic basins are listed in the table below.

Detailed data for individual streams are shown in Figure 9 and Table 2, and annual variations on unimpaired runoff since 1921 for eight streams are shown in Figure 10.

UNIMPAIRED RUNOFF BY AREA

HYDROLOGIC BASIN	WATER YEAR UNIMPAI FLOW IN PERCENT OF NORMAL					
S	1977	1978				
North Coast	14	146				
San Francisco Bay	1	167				
Central Coast	5	295				
South Coastal Area (1)	26	364				
Central Valley Area						
Sacramento San Joaquin and	20	142				
Tulare Lake	20	184				
Lahontan Area (2)	30	128				
ENTIRE STATE	24	158				

⁽¹⁾ Includes Los Angeles, Santa Ana, and San Diego Basins.

⁽²⁾ Includes North and South Lahontan Basins.

FIGURE 9. UNIMPAIRED RUNOFF, 1977-78 WATER YEAR OCTOBER 1 – SEPTEMBER 30




TABLE 2. STREAMFLOW DATA FOR SELECTED STREAMS (METRIC UNITS)

		SNOWMELT I APRIL 1, 1978 -	PER 100 JULY 31, 1978		WATER YEAR OCTOBER 1, 1977 - SEPTEMBER 30, 1978					
BASIN, STREAM, AND STATION	MEASURED FLOW	UNIHPA	IREO RUNOFF		MEASURED FLOW		AIREO RUNOFF (
bishing single s	CUBIC	50-YR AVG (3) CUBIC	PERIOD TOTAL CUBIC	PERCENT	CUBIC	50-YR AVG (3) CUBIC	ANNUAL TOTAL CUBIC	PERCENT		
(1)	OEKAMETRES	OEKAMETRES	DEKAMETRES	AVERAGE	DEKAMETRES	DEKAMETRES	DEKAHETRES	AVERAGE		
NORTH COAST BASIN										
KLAMATH, COPCO TO ORLEANS (4)					6 565 000	5 460 000	6 846 000	125		
SALHON AT SOMESBAR			1 018 000		1 781 000	1 510 000	1 781 000	118		
TRINITY AT LEWISTON EFL AT SCOTIA	90 000	761 000		134	212 000 9 838 000	1 520 000 6 640 000	2 527 000 10 081 000	166 152		
RUSSIAN AT HEALOSBURG					2 408 000	986 000	2 276 000	231		
SAN FRANCISCO BAY BASIN										
NAPA NEAR ST. HELENA					138 000	82 000	138 000	169		
COYOTE CREEK NEAR MADRONE					31 000	56 000	92 000	164		
CENTRAL COAST BASIN										
ARROYO SECO NEAR SOLEDAD NACIMIENTO BELOW NACIMIENTO DAM,					340 000	134 000	341 000	253		
NEAR BRADLEY SANTA YNEZ ABOVE GIBRALTAR GAM,					392 000	236 000	651 000	276		
NEAR SANTA BARBARA					(11)	50 000	247 000	490		
LOS ANGELES BASIN										
SESPE CREEK NEAR FILLMORE					489 000	95 000	489 000	515		
SANTA ANA BASIN						4				
SANTA ANA NEAR HENTONE					141 000	68 000	176 000	259		
SAN DIEGO BASIN SAN LUIS REY AT OCEANSIDE					(11)	47 000	99 000	208		
SACRAHENTO BASIN						(7/2 222	0.660.000	16.0		
INFLOW TO SHASTA (8) SACRAMENTO ABOVE BEND BRIDGE,	2 540 000	2 196 000	2 540 000	116	9 669 000	6 760 000	9 669 000	143		
NEAR RED BLUFF (9)	3 523 000	2 985 000	3 702 000	124	11 123 000	9 810 000	14 806 000	151		
FEATHER, INFLOW TO OROVILLE YUBA AT SMARTVILLE (10)	1 210 000 881 000	2 294 000 1 332 000	2 635 000 1 655 000	115	2 825 000 2 351 000	5 290 000 2 800 000	6 943 000 3 665 000	131		
AMERICAN, INFLOW TO FOLSOM	1 318 000	1 628 000	2 025 000	124	2 877 000	3 170 000	3 953 000	125		
STONY CREEK BELOW BLACK BUTTE DAH					2 877 000	3 170 000	765 000	160		
CACHE CREEK NEAR CAPAY PUTAH CREEK NEAR WINTERS					780 000 (11)	635 000 444 000	1 078 000 752 000	170 169		
					(11)	444 000	/52 000	109		
SAN JOAQUIN BASIN COSUMNES AT MICHIGAN BAR	238 000	163 000	242 000	149	547 000	433 000	600 000	139		
MOKELUHNE, INFLOW TO PARCEE	602 000	575 000	808 000	141	1 042 000	870 000	1 196 000	138		
STANISLAUS, INFLOW TO MELONES	666 000	884 000	1 281 000	145	1 016 000	1 340 000	1 919 000	143		
TUOLUHNE, INFLOW TO DON PEORO	825 000 834 000	1 468 000	2 222 000 1 458 000	151 194	1 180 000 1 283 000	2 210 000 1 130 000	3 327 000 2 164 000	151 190		
MERCEO, INFLOW TO EXCHEQUER ORESTIMBA CREEK NEAR NEWMAN		750 000	1 450 000		(11)	13 000	21 000	157		
SAN JOAQUIN, INFLOW TO HILLERTON	2 290 000	1 468 000	2 876 000	196	3 752 000	2 050 000	4 196 000	205		
TULARE LAKE BASIN										
KINGS, INFLOW TO PINE FLAT	3 220 000	197 000	2 900 000	203	3 836 000	1 940 000	4 112 000	212		
KAWEAN, INFLOW TO TERHINUS LOS GATOS CREEK NEAR COALINGA	669 000	333 000	669 000	201	1 016 000	497 000 4 000	1 027 000 9 000	207 250		
TULE, INFLOW TO SUCCESS	158 000	73 000	158 000	216	312 000	164 000	337 000	205		
KERN, INFLOW TO ISABELLA	1 311 000	518 000	1 311 000	253	1 426 000	773 000	1 885 000	244		
NORTH LAHONTAN BASIN										
SUSAN AT SUSANVILLE	217 000	226 000	297 000		(11)	62 000 470 000	83 000 543 000	135 116		
TRUCKEE, TAHOE TO FARAO (4) WEST FORK CARSON AT WOODFORDS	317 000 76 000	326 000 63 000	387 000 76 000	119	501 000 92 000	86 000	92 000	106		
EAST FORK CARSON NEAR GARONERVILLE	263 000	225 000	263 000	117	339 000	306 000	339 000	392		
WEST WALKER BELOW LITTLE WALKER,	-	•	•							
NEAR COLEVILLE EAST WALKER NEAR BRIOGEPORT	242 000 93 000	176 000 74 000	242 000 116 000	137 155	296 000 139 000	218 000 131 000	296 000 176 000	136 135		
SOUTH LAHONTAN BASIN										
OWENS BELOW LONG VALLEY CAM	933 000	73 000	123 000	167	219 000	175 000	228 000	130		
HOJAVE AT BARSTOW					(11)	1 111 000	278 000	250		
COLORADO RIVER BASIN	8 414 000							102		
COLORADO, INFLOW TO LAKE POWELL		9 424 000	11 095 000	118	12 037 000	13 900 000	14 191 000			

⁽¹⁾ RESERVOIR INFLOW OATA ARE BASED ON OBSERVEO FLOWS AT STATIONS COWNSTREAM FROM LISTED FACILITY.

(2) THE UNIMPAIRED RUNDER OF A STREAM AT ANY STATION IS THE RUNDER WHICH MOULD HAVE OCCURRED UNDER NATURAL CONDITIONS, UNALTERED BY UPSTREAM DIVERSIONS, STORAGE OFFECTIONS, OR BY EXPORTATION OR IMPORTATION OF WATER TO OR FROM OTHER WATERSHEDS.

(3) AVERAGES ARE COMPUTED FOR THE 50-YEAR PERIOD 1921-70.

(4) ACCRETIONS BETWEEN STATIONS.

(5) ESTIMATED VALUE.

(6) INCLUDES FILLIONE BRIGATION COMPANY CANAL.

(7) INCLUDES SOUTHERN CALIFORNIA EDISON COMPANY CANAL.

⁽S) ESTIMATEO VALUE.

(6) INCLUDES SOUTHERN CALIFORNIA EDISON COMPANY CANAL.

(7) INCLUDES SOUTHERN CALIFORNIA EDISON COMPANY CANAL.

(8) COMPUTE DROM OPERATION RECORDS — UMADJUSTED FOR UPSTREAM REGULATION.

(9) UNIMPAINED FLOWS COMPATIBLE TO THOSE AT DISCONTINUED STATION NEAR RED BLUFF.

(10) INCLUDES DEER CREEK.

(11) DATA NOT AVAILABLE AT TIME OF PURLICATION.

TABLE 2. STREAMFLOW DATA FOR SELECTED STREAMS (ENGLISH UNITS)

		SNOWMELT APRIL 1, 1978 -	JULY 31, 1978	WATER YEAR OCTOBER 1, 1977 - SEPTEMBER 30, 1978					
CASIN CTREAM AND STATION	MEASUREO			2)	MEASURED		IREO RUNOFF (2		
BASIN, STREAM, AND STATION	FLOW	50-YEAR	PER 100	PERCENT	FLOW	50-YEAR	ANNUAL	PERCE	
(1)	ACRE-FEET	AVERAGE AC-FT (3)	TUTAL ACRE-FEET	OF AVERAGE	ACRE-FEET	AVERAGE AC-FT (3)	TOTAL ACRE-FEET	OF A V ERA	
ORTH COAST BASIN									
KLAMATH, COPCO TO ORLEANS (4)					5,322,000	4,430,000	5,550,000	125	
SALMON AT SOMESBAR					1,444,000	1,220,000	1,444,000	118	
TRINITY AT LEWISTON	73,000	617,000	825,000	134	172,000	1,230,000	2,049,000	166	
EEL AT SCOTIA					7,976,000	5,380,000	8,173,000	152	
RUSSIAN NEAR HEALOSBURG					1,953,000	799,000	1,845,000	231	
AN FRANCISCO BAY BASIN NAPA NEAR ST. HELENA					112,000	66,500	112,000	169	
COYOTE CREEK NEAR MAGRONE					25,000	45,400	74,000	164	
ENTRAL COAST BASIN					276,000	109,000	276,000	253	
ARROYO SECO NEAR SOLEDAO NACIMIENTO BELOW NACIMIENTO DAM,									
NEAR BRAOLEY SANTA YNEZ ABOVE GIBRALTAR DAM,					318,000	191,000	528,000	276	
NEAR SANTA BARBARA					(11)	40,000	200,000(5)	490(
OS ANGELES BASIN SESPE CREEK NEAR FILLMORE					396,000	76,000	396,000	515	
ANTA ANA BASIN SANTA ANA NEAR MENTONE					114,000	55,000	143,000	259	
AN DIEGO BASIN									
SAN LUIS REY AT OCEANSTOE					(11)	38,200	80,000(5)	208	
ACRAMENTO BASIN INFLOW TO SHASTA (8)	2,059,000	1,780,000	2,059,000	116	7,838,000	5,480,000	7,838,000	143	
SACRAMENTO ABOVE BEND BRIDGE,									
NEAR RED BLUFF (9)	2,856,000	2,420,000	3,002,000	124	9,018,000	7.950,000	12,003,000	151	
FEATHER, INFLOW TO OROVILLE	981,000	1,860,000	2,136,000	115	2,290,000	4,290,000 2,270,000	5,628,000 2,971,000	131	
YUBA AT SMARTVILLE (10)	715,000	1,080,000	1,341,000	124	1,906,000	2,570,000	3,205,000	125	
AMERICAN, INFLOW TO FOLSOM STONY CREEK BELOW BLACK BUTTE GAM	1,000,000		.,0-2,000		(11)	387,000	620,000	160	
EACHE CREEK NEAR CAPAY					632,000	515,000	874,000	170	
PUTAH CREEK NEAR WINTERS					(11)	360,000	610,000	169	
AN JOAQUIN BASIN	103 000	132,000	196,000	149	443,000	351,000	486,000	139	
COSUMNES AT MICHIGAN BAR	193,000 488,000	466,000	655,000	141	845,000	705,000	970,000	138	
MOKELUMNE, INFLOW TO PARDEE STANISLAUS, INFLOW TO MELONES	540,000	717,000	1,039,000	145	824,000	1,090,000	1,556,000	143	
TUOLUMNE, INFLOW TO DON PEDRO	669,000	1,190,000	1,801,000	151	956,000	1,790,000	2,697,000	151	
MERCED, INFLOW TO EXCHEQUER	676,000	608,000	1,182,000	194	1.040.000	920,000	1,755,000	190	
OREST IMBA CREEK NEAR NEWMAN					(11)	10,800	17,000	157	
SAN JOAQUIN, INFLOW TO MILLERTON	1,856,000	1,190,000	2,332,000	196	3,042,000	1,660,000	3,402,000	205	
ULARE LAKE BASIN KINGS, INFLOW TO PINE FLAT	2,611,000	1,160,000	2,351,000	203	3,110,000	1,570,000	3,334,000	212	
KAWEAH, INFLOW TO TERMINUS	542,000	270,000	542,000	201	824,000	403,000	833,000	207	
LOS GATOS EREEK NEAR COALINGA					(11)	3,000	7,500	250	
TULE, INFLOW TO SUCCESS	128,000	59,200	128,000	216	253,000	133,000	273,000	205	
KERN, INFLOW TO ISABELLA	1,063,000	420,000	1,063,000	253	1,156,000	627,000	1,528,000	244	
ORTH LAHONTAN BASIN SUSAN AT SUSANVILLE					(11)	50,000(5)	68,000	135	
TRUCKEE, TAHOE TO FARAD (4)	257,000	264,000	314,000	119	406,000	381,000	440,000	116	
WEST FORK CARSON AT WOODFORDS	61,000	51,100	61,000	120	75,000	70,100	75,000	106	
EAST FORK CARSON NEAR GARDNERVILLE	213,000	182,000	213,000	117	275,000	248,000	275,000	392	
WEST WALKER BELOW LITTLE WALKER,					-1.	177 000	21.0 222	136	
NEAR COLEVILLE EAST WALKER NEAR BRIDGEPORT	196,000 76,000	143,000 60,300	196,000 94,000	137 155	240,000 113,000	177,000	240,000 143,000	136 135	
OUTH LAHONTAN BASIN							105.000		
OWENS BELOW LONG VALLEY DAM	76,000 	59,500	100,000	167	177,000 (11)	142,000 90,000(5)	185,000 225,000	130 250	
OLORADO RIVER BASIN	6,821,000	7,640,000	8,995,000	118	9,758,000	11.300.000	11,504,000	102	
COLORADO, INFLOW TO LAKE POWELL									

⁽¹⁾ RESERVOIR INFLOW DATA ARE BASED ON OBSERVED FLOWS AT STATIONS DOWNSTREAM FROM LISTED FACILITY.

(2) THE UNIMPAIRED RUNOFF OF A STREAM AT ANY STATION IS THE RUNOFF WHICH MOULD HAVE OCCURRED UNDER NATURAL CONDITIONS, UNALTERED BY UPSTREAM DIVENSIONS, STORAGE DEVELOPMENTS, OR BY EXPORTATION OR IMPORTATION OF MATER TO OR FROM OTHER WATERSHEDS.

(3) ACCURATE OF STREAM STATIONS.

(5) ESTIMATED WALL STATIONS.


(6) INCLUDES FILLANCE BRIGATION COMPANY CANAL.

(7) INCLUDES SOUTHERN CALIFORNIA COISON COMPANY CANAL.

(8) COMPUTED FROM OPERATION RECORDS — UNADJUSTED FOR OPSTREAM REGULATION.

(9) UNIMPAIRED FLOWS COMPATIBLE TO THOSE AT DISCONTINUED STATION NEAR RED BLUFF.

(11) DATA NOT AVAILABLE AT TIME OF PUBLICATION.






TABLE 3. STORAGE IN MAJOR RESERVOIRS (METRIC UNITS)

HYDROLOGIC BASIN			CAPAC ITY	STORAGE AS OF OCTOBER 1 CUBIC DEKAMETRES (1)					
AND STREAM	R ES ER VOIR	OPERATOR	CUBIC DEKAMETRES (1)	10-YEAR AVERAGE 1968-1977	1977	1978	DF AVERA		
IDRTH COAST BASIN									
KLAMATH RIVER	UPPER KLAMATH(2)	US BUREAU RECLAMATION	720	363	276	363	100		
KLAMATH RIVER	CLEAR LAKE(2)	US BUREAU RECLAMATION	650	304	180	201	66		
TRINITY RIVER	CLAIR ENGLE	US BUREAU RECLAMATION	3 020	2 081	300	2 307	111		
RUSSIAN RIVER	LAKE MENDOCIND	US CORPS DF ENGINEERS	152	62	16	18	134		
CALAVERAS CREEK	CALAVERAS(3)	CITY-CO SAN FRANCISCO	123	63	33	75	119		
CENTRAL COAST BASIN									
SAN ANTONIO RIVER	SAN ANTONID	MONTEREY CD FCWCD	432	26D	64	291 286	112		
NACINIENTO RIVER SANTA YNEZ RIVER	NAC IMIENTO CACHUMA	MONTEREY CO FCWCD US BUREAU RECLAMATION	432 253	123 200	27 138	238	233 119		
SOUTH COAST (8)									
COYOTE CREEK	CASITAS	CASITAS MUNICIPAL WD	313	244	225	301	123		
PIRU CREEK	LAKE PIRU	UNITED WATER CON DIST	125	28	17	86	300		
PIRU CREEK	PYRAMID(3)	CALIF DEPT WATER RES	211	202(6)	203	206	101		
CASTAIC CREEK	CASTAIC(3)	CALIF DEPT WATER RES	400	72 (6)	72	398	199		
	PERRIS(3)	CALIF DEPT WATER RES	163	93 (6)	93	162	151		
TRIB CAJALCD CREEK	LAKE MATHEWS (4)	METROPOLITAN WATER DIST	224	134	134	101	78		
SAN JACINTO RIVER	LAKE ELSINORE	CALIF DEPT PARKS AND REC	154	27(6)	10	61	227		
SAN LUIS REY RIVER SAN DIEGO RIVER	HENSHAW EL CAPITAN(3)	VISTA IRRIGATION DIST CITY OF SAN DIEGO	252 144	7 2 7	1 19	5 72	61 268		
ENTRAL VALLEY (9)									
SACRAMENTO RIVER	SHASTA	US BUREAU RECLAMATION	5 61D	3 534	778	4 228	120		
CLEAR CREEK	WHISKEYTOWN	US BUREAU RECLAMATION	297	273	268	284	104		
N FK FEATHER RIVER	LAKE ALMANOR	PAC GAS AND ELEC CO	1 620	974	669	1 102	113		
BUCKS CREEK	BUCKS LAKE	PAC GAS AND ELEC CO	127	74	46	89	12D		
FEATHER RIVER	DRDVILLE	CALIF DEPT WATER RES	4 370	2 846(6)	1 129	3 385	119		
NDRTH YUBA RIVER	NEW BULLARDS BAR	YUBA CD WATER AGENCY	1 190	687(6)	318	718	104		
SOUTH YUBA RIVER	SPAULDING SYSTEM	PAC GAS AND ELEC CD	169	79	76	78	98		
BEAR RIVER	CAMP FAR WEST	SO SUTTER WATER DIST	127	53	4	83	156		
M FK AMERICAN RIVER	FRENCH MEADOWS	PLACER CO WATER AGENCY	165	96	47	120	124		
RUBICON RIVER	HELL HOLE	PLACER CO WATER AGENCY	257	150	96 70	181 250	120		
SILVER CREEK	UNION VALLEY	SACRAMENTO MUN UD	334	189	181	863	133 115		
AMERICAN RIVER	FDLSDM	US BUREAU RECLAMATION	1 250	754	101	47	141		
STONY CREEK	BLACK BUTTE	US CORPS OF ENGINEERS	197	33 70	0	105	149		
CACHE CREEK	CLEAR LAKE	YOLO CDUNTY FCWCD YOLO CDUNTY FCWCD	518 370	0(6)	0	194	0		
N FK CACHE CREEK PUTAH CREEK	INDIAN VALLEY	US BUREAU RECLAMATION	1 970	1 567	936	1 383	88		
N KF MOKELUMNE RIVER	SALT SPRINGS	PAC GAS AND ELEC CO	171	83	5	111	135		
MOKELUMNE RIVER	PARDEE	EAST BAY MUN UD	259	216	102	218	101		
MOKELUMNE RIVER	CAMANCHE	EAST BAY MUN UD	533	321	68	417	13D		
CALAVERAS RIVER	NEW HDGAN	US CORPS OF ENGINEERS	401	157	14	200	128		
STANISLAUS RIVER	MELDNES	PAC GAS AND ELEC CO	139	12	4	64	534		
CHERRY CREEK	CHERRY LAKE	CITY-CO SAN FRANCISCO	332	173	130	242	139		
TUOLUMNE RIVER	HETCH HETCHY	CITY-CD SAN FRANCISCO	444	270	139	402	149		
TUDLUMNE RIVER	DDN PEDRD	TURLOCK-MODESTD 1D	2 500	1 003(6)	379	1 943	193		
MERCED RIVER	LAKE MCCLURE	MERCED IRRIG DISTRICT	1 270	616	116	945	153		
SAN JOAQUIN RIVER	MAMMOTH POOL	SD CALIFORNIA EDISON CO	152	41	25	101	246		
MOND CREEK	THDMAS A EDISON	SO CALIFORNIA EDISON CO	154	93	7	153	164		
STEVENSON CREEK	SHAVER LAKE	SD CALIFORNIA EDISON CD	167	80	36	158	197		
SAN JOAQUIN RIVER	MILLERTON LAKE	US BUREAU RECLAMATION	643	218	243	466	214		
SAN LUIS CREEK	SAN LUIS (3)	US BUREAU REC-CALIF DWR	2 52D	1 706(6)	338	2 120	123		
HELMS CREEK	C DURTR I GHT	PAC GAS AND ELEC CO	152	43	1	88	205		
N FK KINGS RIVER	WISHON	PAC GAS AND ELEC CO	158	107	99	123	115		
KINGS RIVER	PINE FLAT	US CORPS OF ENGINEERS	1 230	454	84	909	200 131		
KAWEAH RIVER KERN RIVER	TERMINUS ISA BELLA	US CORPS OF ENGINEERS US CORPS OF ENGINEERS	185 703	21 199	12 44	27 454	229		
AHONTAN (10)									
LITTLE TRUCKEE RIVER	STAMPEDE (2)	US BUREAU RECLAMATION	28 D	150(6)	38	76	5D		
TRUCKEE RIVER OWENS RIVER	LAKE TAHOE(2,7) LAKE CROWLEY	US BUREAU RECLAMATION	919 227	575 148	0 64	220 212	38 143		
	LANE UNUWLET	LDS ANGELES DEPT WP	22/	140	04	*1*	. 45		
COLORADO RIVER BASIN	LAKE POWELL (2,7)	US BUREAU RECLAMATION	30 800	18 335	19 914	20 430	111		
COLORADO RIVER	LAKE MEAD(2,7)	US BUREAU RECLAMATION	32 20D	22 476	24 923	25 742	115		
CDLORADO RIVER	LAKE MOHAVE(2,7)	US BUREAU RECLAMATION	2 230	1 779	1 807	1 831	103		
COLORADO RIVER	LAKE HAVASU(2,7)	US BUREAU RECLAMATION	764	696	698	699	101		

CAPACITY AND STDRAGE VALUES RDUNDED TO NEAREST THREE SIGNIFICANT NUMBERS.
INTERSTATE RESERVOIR USED JOINTLY BY CALIFORNIA AND ADJACENT STATES.
INCLUDES FOREION WATER.
STORES ONLY IMPORTED COLORADD RIVER WATER.
NEW RESERVOIR -- AVERAGE CONSIDERED EQUAL TO CURRENT STORAGE.
LESS THAN 10-YEAR AVERAGE.
DATA BASED DA ACTIVE OR USABLE CAPACITY TABLES.
SOUTH COAST INCLUDES LOS ANGELES, SANTA ANA, AND SAN DIEGO BASINS.
CENTRAL VALLEY INCLUDES SACRAMENTO, SAN JOAQUIN, AND TULARE LAKE BASINS.
LAHONTAN INCLUDES NORTH AND SOUTH LAHONTAN BASINS.

⁽³⁾ (4) (5) (6) (7)

TABLE 3. STORAGE IN MAJOR RESERVOIRS (ENGLISH UNITS)

HYDROLOGIC BASIN			CAPACITY 1000	STORAGE AS OF OCTOBER 1 1000 ACRE-FEET (1)					
AND STREAM	RESERVOIR	OPERATOR	ACRE-FEET	10-YEAR AVERAGE 1968-1977	1977	1978	PERCENT OF AVERA		
RTH COAST BASIN									
KLAMATH RIVER	UPPER KLAMATH(2)	US BUREAU RECLAMATION	584	294	224	294	100		
KLAMATH RIVER	CLEAR LAKE(2)	US BUREAU RECLAMATION	527	247	147	163	66		
TRINITY RIVER	CLAIR ENGLE	US BUREAU RECLAMATION	2,450	1,687	242	1,870	111		
RUSSIAN RIVER	LAKE MENDOCINO	US CORPS OF ENGINEERS	123	50	13	66	134		
N FRANCISCO BAY BASIN CALAVERAS CREEK	CALAVERAS (3)	CITY-CO SAN FRANCISCO	100	51	27	61	119		
NTRAL COAST BASIN									
SAN ANTONIO RIVER	SAN ANTONIO	MONTEREY CO FCWCD	350	211	52	236	112		
NACIHIENTO RIVER	NAC IMIENTO	MONTEREY CO FCWCO	350	100	22	232	233		
SANTA YNEZ RIVER	CACHUMA	US BUREAU RECLAMATION	205	162	112	193	119		
UTH COAST (8)			254	198	182	244	100		
COYOTE CREEK	CASITAS LAKE PIRU	CASITAS MUNICIPAL WO UNITED WATER CON DIST	101	23	14	70	123		
PIRU CREEK PIRU CREEK	PYKAHID(3)	CALIF DEPT WATER RES	171	164(6)	165	167	101		
CASTAIC CREEK	CASTAIC(3)	CALIF DEPT WATER RES	324	162 (6)	58	324	199		
ASTATE CREEK	PERRIS(3)	CALIF DEPT WATER RES	132	86(6)	75	131	151		
RIB CAJALCO CREEK	LAKE MATHEWS (4)	METROPOLITAN WATER DIST	182	105	109	82	78		
AN JACINTO RIVER	LAKE ELSTNORE	CALIF DEPT PARKS AND REC	125	22(6)	8	50	227		
SAN LUIS REY RIVER	HENSHAW	VISTA IRRIGATION DIST	204	6	ĭ	4	61		
SAN DIEGO RIVER	EL CAPITAN(3)	CITY OF SAN DIEGO	117	22	15	58	268		
TRAL VALLEY (9)			1, 550	0.065	(2)	0.100	100		
ACRAHENTO RIVER	SHASTA	US BUREAU RECLAMATION	4,550	2,865	631	3,428	120		
LEAR CREEK	WHISKEYTOWN	US BUREAU RECLAMATION	1,310	221 794	217 542	230	104		
FK FEATHER RIVER	LAKE ALMANOR	PAC GAS AND ELEC CO	1,310	60	542 37	893 72	113		
UCKS CREEK	BUCKS LAKE	PAC GAS AND ELEC CO CALIF DEPT WATER RES	3,540	2,307(6)	37 915	2.744	119		
EATHER RIVER IORTH YUBA RIVER	OROVILLE NEW BULLARDS BAR	YUBA CO WATER AGENCY	961	557(6)	258	582	104		
OUTH YUBA RIVER	SPAULDING SYSTEM	PAC GAS AND ELEC CO	137	64	62	63	98		
EAR RIVER	CAMP FAR WEST	SO SUTTER WATER DIST	103	43	3	67	156		
FK AMERICAN RIVER	FRENCH MEADOWS	PLACER CO WATER AGENCY	134	78	38	97	124		
UBICON RIVER	HELL HOLE	PLACER CO WATER AGENCY	208	122	78	147	120		
SILVER CREEK	UNION VALLEY	SACRAMENTO MUN UO	271	153	57	203	133		
MERICAN RIVER	FOLSOM	US BUREAU RECLAMATION	1,010	611	147	700	115		
STONY CREEK	BLACK BUTTE	US CORPS OF ENGINEERS	160	27	1	38	141		
ACHE CREEK	CLEAR LAKE	YOLO COUNTY FOWCD	420	57	0	85	149		
FK CACHE CREEK	INDIAN VALLEY	YOLO COUNTY FEWED	300	0(6)	0	157	0		
PUTAH CREEK	LAKE BERRYESSA	US BUREAU RECLAMATION	1,600	1,670	759	1,121	88		
N KE HOKELUMNE RIVER	SALT SPRINGS	PAC GAS AND ELEC CO	139	67	4	90	135		
AOKELUMNE RIVER	PARDEE	EAST BAY HUN UD	210	175	83	177	101		
HOKELUMNE RIVER	CAMANCHE	EAST BAY HUN UD	432	260	55	338	130		
ALAVERAS RIVER	NEW HOGAN	US CORPS OF ENGINEERS	325	127	11	162	128		
STANISLAUS RIVER	MELONES	PAC GAS AND ELEC CO	113	10	3	52	534		
HERRY CREEK	CHERRY LAKE	CITY-CO SAN FRANCISCO	269	140	105	196	139 149		
UOLUMNE RIVER	HETCH HETCHY	CITY-CO SAN FRANCISCO	360	219	113 307	326 1,575	149		
UOLUMNE RIVER	DON PEDRO	TURLOCK-HOOESTO ID	2,030 1,030	813 (6) 499	307 94	766	153		
ERCED RIVER AN JOAQUIN RIVER	LAKE MCCLURE MAMMOTH POOL	MERCED IRRIG DISTRICT SO CALIFORNIA EDISON CO	1,030	33	20	82	246		
IAN JOAQUIN RIVER NONO CREEK	THOMAS A EDISON	SO CALIFORNIA EDISON CO	123	75	6	124	164		
TEVENSON CREEK	SHAVER LAKE	SO CALIFORNIA EDISON CO	135	65	29	128	197		
IAN JOAQUIN RIVER	MILLERTON LAKE	US BUREAU RECLAMATION	521	177	197	379	214		
SAN LUIS CREEK	SAN LUIS (3)	US BUREAU REC-CALIF DWR	2,040	1,383(6)	2.74	1,719	123		
IELHS CREEK	COURTRIGHT	PAC GAS AND ELEC CO	123	35	7/1	71	205		
FK KINGS RIVER	WISHON	PAC GAS AND ELEC CO	128	87	80	600	115		
INGS RIVER	PINE FLAT	US CORPS OF ENGINEERS	1,000	368	68	737	200		
AWEAH RIVER	TERMINUS	US CORPS OF ENGINEERS	150	17	10	22	131		
CERN RIVER	ISABELLA	US CORPS OF ENGINEERS	570	161	36	368	229		
HONTAN (10)			22.7	122(6)	21	62	50		
LITTLE TRUCKEE RIVER	STAMPEDE (2)	US BUREAU RECLAMATION	227 745	122 (6) 466	31	178	38		
TRUCKEE RIVER OWENS RIVER	LAKE TAHOE(2,7) LAKE CROWLEY	US BUREAU RECLAMATION LOS ANGELES DEPT WP	184	120	52	172	143		
LORADO RIVER BASIN									
COLORADO RIVER	LAKE POWELL (2,7)	US BUREAU RECLAMATION	25,000	14,864	16,144	16,563	111		
				18,221	20,205	20,860	115		
	LAKE MEAD(2,7)	US BUREAU RECLAMATION	26,100	10,221					
COLORADO RIVER COLORADO RIVER	LAKE MEAD(2,7) LAKE MOHAVE(2,7)	US BUREAU RECLAMATION US BUREAU RECLAMATION US BUREAU RECLAMATION	1,810	1,442	1,465	1,484	103		

CAPACITY AND STORAGE VALUES ROUNGED TO NEAREST THREE SIGNIFICANT NUMBERS.

CAPACITY AND STORAGE VALUES ROUNGED TO MEAREST THREE SIGNIFICANT NUMBERS INTERSTATE RESERVOIR USED JOINTLY BY CALIFORNIA AND ADJACENT STATES. INCLUDES FOREIGN WATER, STORES ONLY IMPORTED COLORADO RIVER WATER.

NEW RESERVOIR -- AVERAGE CONSIDERED EQUAL TO CURRENT STORAGE. LESS THAN 10-YEAR AVERAGE.

LESS THAN 10-YEAR AVERAGE.

SOUTH COAST INCLUDES LOS ANDELES, SANTA ANA, AND SAN DIEGO BASINS. CENTRAL VALLEY INCLUDES SACRAHENTO, SAN JOACHIN, AND TULLARE LAKE BASINS. LAHONTAN INCLUDES NORTH AND SOUTH LAHONTAN BASINS.

⁽⁵⁾ (6) (7) (8) (9) (10)

RESERVOIR STORAGE

On October 1, 1977 water storage in 153 major reservoirs had been reduced by two consecutive years of drought to an all-time low record of 9.74 cubic dekametres (7.9 million acre-feet), or 38 percent of average and 23 percent of available capacity.

Heavy precipitation in the first half of January 1978 brought relief from the drought and caused some minor reservoirs to fill and some major reservoirs to reach normal storage levels by mid month. For example, Folsom Reservoir in the Sacramento River Basin rose rapidly from an all-time low to a level requiring flood control releases.

By June 1, 1978, reservoir storage in the San Joaquin Valley varied from 75 percent of average in the San Joaquin and Kings Rivers to 142 percent of average on the Kern River. Storage in these reservoirs reflected flood control operations space for anticipated above-average inflows from snowmelt. Unusual below-normal temperatures at higher elevations until mid July, however, gave reservoir operators and water managers time to adequately control the season's large runoff.

On September 30, 1978, storage in 153 major reservoirs was 30.3 million dam^3 (24.6 million ac.-ft.) or 127 percent of average and 71 percent of available capacity, an increase of 20.6 million dam^3 (16.7 million ac.-ft.).

TABLE 4. SUMMARY OF RESERVOIR STORAGE DATA
THOUSANDS OF CUBIC DEKAMETRES(THOUSANDS OF ACRE-FEET)

BASIN	NUMBER OF RESERVOIRS	TOTAL CAPAC ITY	10-YEAR AVERAGE 1968-77	STORAGE OCTOBER 1 1978	PERCENT OF AVERAGE	PERCENT OF CAPACITY
INTRASTATE:						
NORTH COAST	8	3 630	2 387	2 667	112	74
		2,940	1,935	2,162		
SAN FRANCISCO BAY	18	859	477	517	1 08	60
		696	387	419		
CENTRAL COAST	9	1 300	714	966	135	75
		1,059	579	783		
SOUTH COAST (1)	31	2 840	1 278	1 871	146	66
		2,300	1,036	1,517		
SACRAMENTO	43	20 100	12 305	14 010	114	70
		16,300	9,976	11,358		7.5
SAN JOAQUIN AND TULARE LA	KE 36	13 200	6 353	9 863	155	75
		10,700	5,150	7,996		0.5
LAHONTAN (2)	8	525	350	444	127	85
		426	284	360		
SUBTOTAL	153	42 400	23 865	30 338	127	71
300101AL	177	34,400	19,347	24,595	,	, .
		54,400	13,377	27,000		
INTERSTATE:						
NORTH COAST	3	1 490	714	464	84	40
		1,210	579	376		
LAHONTAN (2)	5	1 330	783	44	52	31
		1,080	635	36		-1
COLORADO RIVER (3)	4	66 000	43 285	47 342	113	74
		53,500	35,091	38,380		
SUBTOTAL (3)	12	68 800	44 782	47 850	111	72
SOULOUVE (3)	1.6	55,800	35,305	38,792		, -
		55,000	52,505	50,752		
TOTAL (3)	165	111 000	68 647	78 194	114	30
		90,200	55,652	63,387		

⁽¹⁾ INCLUDES LOS ANGELES, SANTA ANA, AND SAN DIEGO BASINS.

⁽²⁾ INCLUDES NORTH AND SOUTH LAHONTAN BASINS
(3) INCLUDES NORTH AND SOUTH LAHONTAN BASINS
(3) INCLUDES DATA FOR LAKE MEAD AND LAKE POWELL WHICH REGULATE FLOW OF THE LOWER COLORADO RIVER, THE MAJOR SOURCE OF WATER FOR THE COLORADO RIVER BASIN AND SOUTH COAST AREA.

WATER SUPPLY FORECAST VERIFICATION

Tabulated below are 1977-1978 water supply forecasts published in Bulletin 120, "Water Conditions in California", and the observed unimpaired runoff for 25 major forecast points. Error percentages compare May 1 forecasts to the observed. Forecasts are always subject to limitations in forecasting procedures, and in the uncertainty of future weather. Forecast errors due to forecasting procedures usually range between 5-10 percent. On April 1, 1978, there was a heavy snowpack at the higher elevations, above 1 980 metres (6,500 feet). Cold temperatures and above-normal precipitation during April further increased the snowpack. Average temperatures for May and June continued cool. Absent were the periods of high temperature which normally cause heavy snowmelt runoff. It was not until mid July that temperatures significantly exceeded normal. Consequently, forecasted April-July snowmelt potential was not realized until late August or early September, thus creating an apparent overforecasted situation.

TABLE 5. COMPARISONS OF WATER SUPPLY FORECASTS WITH OBSERVED UNIMPAIRED RUNOFF

Flows In 1 000 Cubic Oekametres -- 1,000 Acre-feet

STREAM AND STATION				FORE	CASTS				OBSE		FORECAS'	
SIREAR AND STATION	FEBRUAPR-JULY			H 1 WATER YR		WATER YR	MA*			OWS WATER YR	MAY 1 F	
TRINITY AT LEWISTON	1 011 820	2344 1,900	1 110 900	2 467 2,000	1 092 885	2 572 2,085	1 246 1,010	2 732 2,215	1 018 825	2 519 2,050	+ 22	+ 8
SACRAMENTO R INFLOW TO SHASTA LAKE	481	1 523	537 435	1 678 1,360	567 460	1,776	703 570	1 918	491 398	1 708 1,385	+ 43	+ 12
McCLOUD R INFLOW TO SHASTA LAKE	555 450	1 653	580 470	1 813 1,470	617 500	1 881	715 580	1 992 1,615	510 414	1 711	+ 40	+ 16
PIT R INFLOW TO SHASTA LAKE	125 101	4 348 3,525	1 264	4 132 3,350	1 283 1,040	4 225 3,425	1 419 1,150	4 373 3,545	1 356 1,099	4 317 3,500	+ 5	+ 1
TOTAL INFLOW TO SHASTA LAKE	2 368 1,920	8 783 7,120	2 467 2,000	9 153 7,420	2 541 2,060	9 633 7,810	3 084 2,500	10 189 8,260	2 540 2,059	9 668 7,838	+ 21	+ 5
SACRAMENTO R. ABOVE BEND BRIDGE	3 238 2,625	12 828 10,400	3 330 2,700	13 569 11,000	3 454 2,800	14 481 11,740	4 317 3,500	15 406 12,490	3 703 3,002	14 806 12,003	+ 17	+ 4
FEATHER R INFLOW TO OROVILLE RES.	2 874 2,330	6 846 5,550	3 108 2,520	6 883 5,580	3 121 2,530	7 494 6,075	3 639 2,950	8 030 6,510	2 509 2,034	6 942 5,628	+ 45	+ 16
YUBA R. AT SMARTVILLE	1 493	3 300 2,675	1 604	3 306 2,680	1 604	3 690 2,910	1,580	3 947 3,200	1 654	3 665 2,971	+ 18	+ 8
AMERICAN R INFLOW TO FOLSOM RES.	1 856 1,505	3 700 3,000	2 035 1,650	3 750 3,040	2 035 1,650	3 953 3,205	2 516 2,040	3,605	1,641	3 953 3,205	+ 24	+ 12
COSUMNES R. AT MICHIGAN BAR	197	514 417	210 170	506 410	216 175	574 465	296 240	654 530	242 196	6 00 486	+ 22	+ 9
MOKELUMNE R INFLOW TO PARDEE RES.	691 560	1 005 815	759 615	1 086	802 650	1 184 960	937 760	1 326	807 654	1,002	+ 16	+ 1
STANISLAUS R INFLOW TO HELONES RES.	1 061 860	1 499	1 234	1,400	1 357 1,100	2 035 1,650	1 653	2 276 1,845	1 282	1 919	+ 29	+ 19
TUOLUMNE R INFLOW TO OON PEORO RES.	1 789	2 578 2,090	2 159 1,750	3 084 2,500	2 381 1,930	3 441 2,790	2 837 2,300	3 916 3,175	1,801	3 327 2,697	+ 28	+ 18
MERCEO R INFLOW TO EXCHEQUER RES.	987 800	1 456	1 178	1,430	1 357	1 998	1 628 1,320	2 288 1,855	1 458	2 165 1,755	+ 12	+ 6
SAN JOAQUIN R INFLOW TO MILLERTON LAKE	2 097 1,700	2 874	2 553 2,070	3 528 2,860	3 158 2,560	4 286 3,475	3 565 2,890	4 706 3,815	2 877	4 196 3,402	+ 24	+ 11
KINGS R INFLOW TO PINE FLAT RES.	1 974	2 671	2 399	3 293 2,670	2 899 2,350	3 873 3,140	3 207 2,600	3,400	2 900	3,334	+ 11	+ 4
KAWEAH R INFLOW TO TERMINUS RES.	419 340	574 465	537 435	802 650	678 550	999 810	740 600	1 067 865	669 542	1 028 833	+ 5	+ 4
TULE R INFLOW TO SUCCESS RES.	93 .75	168 136	117 95	271 220	148 120	1 321	167	339 275	158	337 273 1 885	+ 17	+ 12
KERN R INFLOW TO ISABELLA RES.	648 525	919 745	1 048 850	1 493	1,200	2 054 1,665	1,240	2 115 1,715	1,063	1,528	+ 19	+ 15
TRUCKEE R. TAHJE TO FARAO	407 330	543 440	419 340	543 440	401 325	555 450	469 380	623 505	392 318	440		
LAKE TAHOE RISE	1.60		1.80		1.75	1.12	1.90	413	1.37	339	+ 39	+ 22
EAST CARSON R. NEAR GARONERVILLE	259 210	327 265	321 260	395 320	333 270	413 335	358 290 105	335 123	213 75	275 93	+ 39	+ 33
WEST CARSON AT WOODFORDS	74 60	93 .75	93	90	93	90	85 160	100	61	176	+ 38	+ 26
EAST WALKER R. NEAR BRIDGEPORT	105 85	142	136	173	142	179	130	180	94 242	143 296	+ 17	+ 13
WEST WALKER R. NEAR COLEVILLE	180	253 205	259 210	308 250	265 215	315 255	284	333 270	196	240		''


FORECAST ERROR IN PERCENT = (FORECAST - OBSERVED)

OBSERVED X 100

MAY I APRIL-JULY FORECAST AVERAGE ABSOLUTE ERROR = 28 % (25 FORECAST)
MAY I WATER YEAR FORECAST AVERAGE ABSOLUTE ERROR = 18 % (18 FORECAST)

Note: Acre - Feet = 1.2335 Cubic Dekometres

FIGURE 11. WATER TRANSFERS AND OUTFLOW, 1977-78



WATER TRANSFERS AND OUTFLOW

Heavy rainfall and near-record snow accumulation during the 1977-78 water year brought relief to California following two years of drought. Runoff and reservoir storage reached levels of near normal or greater in almost all portions of the State.

State Water Project diversions from the Delta increased from 936,000 cubic dekametres (759,000 acre-feet) in 1976-77 to 2 499 000 dam 3 (2,026,000 ac.-ft.) in 1977-78. A large portion of this, 1 056 000 dam 3 (856,000 ac.-ft.) was used to increase the depleted storage in San Luis Reservoir. State Water Project deliveries to Southern California increased from 222 000 dam 3 (180,000 ac.-ft.) to 614 000 dam 3 (498,000 ac.-ft.).

Central Valley Project Delta diversions also greatly increased over the previous year from 1 788 000 dam 3 (1,450,000 ac.-ft.) to 2 975 000 dam 3 (2,412,000 ac.-ft.). Of this amount 727 000 dam 3 (589,000 ac.-ft.) was put into storage at San Luis Reservoir. Trinity Rîver dîversions dropped sharply from 1 626 000 dam 3 (1,318,000 ac.-ft.) to 269 000 dam 3 (218,000 ac.-ft.).

Hetch Hetchy Aqueduct diversions for San Francisco decreased from 317 000 dam³ (257,000 ac.-ft.) to 175 000 dam³ (142,000 ac.-ft.). Emergency Delta diversions to supply water for the drought stricken San Francisco Bay area ended in January 1978 as winter rains eased the critical shortage.

Los Angeles Aqueduct diversions increased from 421 000 dam^3 (341,000 ac-ft.) in 1976-77 to 522 000 dam^3 (423,000 ac.-ft.) in 1977-78.

Colorado River Aqueduct diversions decreased from 1 335 000 $\rm dam^3$ (1,082,000 ac.-ft.) to 1 100 000 $\rm dam^3$ (892,000 ac.-ft.) reflecting the increased amount received from the State Water Project.

Klamath River inflow from Oregon increased from 1 134 000 $\rm dam^3$ (919,000 ac.-ft.) in 1976-77 to 2 019 000 $\rm dam^3$ (1,637,000 ac.-ft.) in 1977-78. The estimated outflow of the Truckee, Carson, and Walker Rivers to Nevada increased from 512 000 $\rm dam^3$ (415,000 ac.-ft.) to 1 067 000 $\rm dam^3$ (1,108,-00 ac.-ft.).



Tow truck operator clings to safety rope in Santa Ana River after attaching line to a trapped truck, he was unharmed





A man and woman comfort one another in the Upper Big Tujunga Canyon area. They were rescued by helicopter after a dam cellapsed, sending a wall of water through a trailer park.



Red Hill Country Club Drive in Rancho Cucamonga after water had subsided



San Bernardino sheriff's helicopter participates in search for survivors from a car in Cajon Creek Wash. None were found

CHAPTER III - FLOOD EVENTS

The sudden turnaround of meteorologic events experienced in water year 1977-78 is convincing evidence that the variability of California weather is its greatest consistency. We are reminded, too, that the transition between drought and flooding can be abrupt and dramatic.

Precipitation during the first two months of water year 1977-78 beginning October 1 was a disappointment to most Californians, who were still feeling the effects of two consecutive years of withering drought. The series of weather fronts that moved through Northern California in late October brought substantial precipitation to some North Coast stations but only light amounts to the northern and central Sierra basins. The remainder of the State remained virtually dry. The estimated 130 percent of normal precipitation needed to alleviate the critical effects of the drought appeared to be even more remote following the meager rainfall of the fall months. Although additional weather fronts passed through Northern California during November, bringing seasonal totals to near normal in some areas on the North Coast, the water supply outlook for the rest of the State at the onset of the winter quarter remained bleak.

And then the rains came! In mid-December, the high-pressure ridge that had persisted off the California coast since November 1975 was dislodged, and the long-awaited storms swept across parched California. Heavy rains fell in the valley and foothill areas, and deep snows blanketed the mountains. Particularly heavy rains occurred in portions of Southern California, where 102 to 152 mm (4 to 6 in.) fell in a four-day period, causing wind and rain damage and numerous mudslides. By the end of December, rainfall in all of California was above or near normal, and storage in the 143 major reservoirs had risen to 50 percent of the 10-year average. The drought cycle was beginning to weaken.

Steady rains in lighter amounts continued for the first twenty days of January 1978, raising seasonal totals to considerably above normal; yet, because of the extremely parched condition of the soils, runoff was not proportionate. Three of the fixed weirs on the Sacramento River, however, overflowed for the first time since 1975, and local flooding occurred in many communities from Tehama on the Upper Sacramento River to San Diego.

From January 20 through February 5, California welcomed some relief from the drenching. Then, on February 6, the downpour began again for another ten days, resulting in swollen rivers, flood stages, mudslides, and loss of life. Southern California suffered significant damage.

Another onslaught of storms hit the State during the final days of February and extended well into March. Again, the major impact of the storms was felt in Southern California, where persistent torrential rains triggered more mudslides and floodflows.

By March 8, 1978, thirteen counties were declared federal disaster areas. Kern, Kings, Los Angeles, Monterey, Orange, Riverside, San Bernardino, Santa Barbara, Tulare, Ventura, Inyo, San Diego, and San Luis Obispo became eligible for financial assistance under the Disaster Relief Act of 1974, Public Law 93-288.

The wet weather pattern continued through April, extending the string of consecutive months of well-above-normal precipitation to five. On May 1, the California Cooperative Snow Surveys reported seasonal statewide precipitation to that date at 155 percent of normal and major reservoir storage at 105 percent of the 10-year average (twice the amount of the previous year). Californians were now assured of ample water supplies for the remainder of 1978.

A summary of events within pertinent hydrographic areas follows:

North Coastal Hydrographic Area

Four significant storms affected this area during the 1977-78 water year: November 22-26, December 15, January 6-20, and February 3-14. The impact of the first major storm of the season during late November was confined primarily to the Smith River Basin. Later weather fronts, December-February, tracked further south, generating flood and warning stages on most rivers and streams; however, flood damage throughout the area was minimal.

San Francisco Bay Hydrographic Area

On January 16 the crest of the Napa River at St. Helena rose to $1.0~\mathrm{m}$ (3.2 ft.) above the flood stage, but damage at St. Helena was limited to levee erosion. By the time the waters reached Napa, the flows were below warning stages.

Later in the month, a combination of high tides and high flows caused levee breaches on the Napa River in Vallejo. Water spread over an area of about $5.2~\rm km^2$ (2 mi. 2), and some houses and mobile homes were flooded.

Central Coastal Hydrographic Area

The Salinas River rose above flood stage five times between January 16 and March 5. On January 10 it crested at 5.67 m (18,6 ft.) at Bradley, with a discharge of 2 000 m 3 /s (70,600 ft. 3 /s). The flood stage at Bradley is 3.35 m (11.0 ft.). Although flood control was provided by both the Nacimiento and San Antonio dams, high flows inundated about 8 100 ha (20,000 ac.) of farmland along the Salinas River. Damage from this flooding, though not as great as the 1969 flood, was estimated at \$7 million.

During this period, numerous roads were closed, and the Carmel River channel was heavily eroded.

Central Valley Hydrographic Area

Sacramento River Basin

From January 8 to 19, a number of mobile homes located in the Sacramento River flood plain between Red Bluff and Vina had to be evacuated. During this same period, a number of additional flood-related incidents took place in the Sacramento River Basin.

A levee failed on the southwest side of Grizzly Island in the Delta, but little damage resulted because the island primarily comprises marshlands used by hunting clubs. In another portion of the Delta, Department of Water Resources crews were sent to Veale Tract to contend with flooding that resulted from high tides and heavy rainfall runoff.

Department of Water Resources crews also placed canvas on 700 metres (2,300 feet) of levee slope in the Cache Creek Settling Basin to retard wavewash erosion. During this period, high flows damaged a levee on Murphy Slough in Maintenance Area 15, southwest of Chico, to the extent that repairs required 7 600 m 3 (10,000 yd. 3) of earth and 180 tonne (200 tons) of rock.

In February, high winds and high tides threatened to overtop Bradford Island levees in the Sacramento-San Joaquin Delta. Crews from the Forestry Conservation Camp, directed by DWR personnel, sandbagged low spots on the levees to protect the island from flooding.

From March 4 through March 19, the fixed weirs on the Sacramento River overflowed continuously, inundating prime agricultural lands; however, these flows had little effect on planting programs.

San Joaquin River Basin

Between February 1 and May 31, 4 300 000 cubic dekametres (3,500,000 acre-feet), or 198 percent of normal flows, passed through the lower San Joaquin River. In an effort to reduce the flows, local water agencies diverted as much water as possible for preirrigation and ground water recharge. The U. S. Bureau of Reclamation delivered Friant Dam water to contractors along the Friant-Kern Canal and the Madera Canal, much of which was recharged to ground water. Additional water was diverted to the Madera Canal and released down the Chowchilla River, Ash Slough, and Berenda Slough to recharge the ground water through these permeable channels.

In spite of these efforts, the San Joaquin River at Newman reached or exceeded the warning stage of 19.2 metres (63.0 feet) from February 15 through February 19; March 9 through March 14; and April 4 through May 11. The San Joaquin River flows were above the warning stage at Vernalis from April 8 through May 15. During this last period, the river flow fluctuated from 680 cubic metres per second (24,000 ft. 3/s) to near 1 080 m³/s (38,000 ft. 3/s), and trailer parks and resort areas located within the flood plain near Vernalis, Mossdale, and Maze Road Bridge were evacuated.

By May 1, the high flows of $140~\mathrm{m}^3/\mathrm{s}$ (4,900 ft. $^3/\mathrm{s}$) in Kings River North that were being diverted away from Tulare Lake were rapidly eroding the right bank levee at points 10.5 to 12 kilometres (6.5 to 7.5 miles) downstream from Crescent Weir. The Kings River Conservation District rushed 2 700 tonne (3,000 ton) of concrete rubble to the erosion sites, and on May 3 the Department of Water Resources declared a preemergency and assigned trained personnel to assist in patrolling the levee. By May 4, the U. S. Corps of Engineers began placing rock revetment along the damage area, and, by May 15, seven sites had been revetted with 6 350 tonne (7,000 ton) of rock. The 24-hour levee patrolling program initiated on May 2 was terminated.

Tulare Lake Basin

This season's rainfall on the Tulare Lake bed area was about 381 millimetres (15 inches), or double the normal amount. This caused exceptionally heavy local runoff to the low portion of the Lake, and by mid-March more than 16 000 hectares (41,000 acres) were inundated to an average depth of 0.6 metres (2 feet). Much of this water was runoff from Deer Creek, Poso Creek, Avenal Gap, and the White River.

With the Kings, Kaweah, Tule and Kern Rivers promising continued heavy flows, several methods were devised to stop or effectively prevent their flows from reaching the lakebed, as described in the following paragraphs.

Kings River Flows

Kings River flows were diverted to Kings River North (Fresno Slough) and into the San Joaquin River. Also, between March 29 and April 11, 1978, 20 temporary pumps installed at weirs along Kings River South moved 3 200 cubic dekametres (2,600 ac.-ft.) of water north (upstream) from Tulare Lake and into Fresno Slough.

Kaweah River Flows

To the extent possible, the 1 200 kilometres (750 miles) of rivers, streams, and canals within the Kaweah River system were used to impound water enabling considerable amounts to percolate into ground water aquifers. In addition, water was taken from the river and spread over 2 000 hectares (5,000 acres) of percolation basins. An additional 11 200 dam³ (9,100 ac.-ft.) was pumped into the Friant-Kern Canal and released into the Kern River. In spite of this, 20 900 dam³ (17,000 ac.-ft.) of unwanted water made its way to Tulare Lake through the Kaweah River system during February through May.

Tule River Flows

Facilities to make use of the runoff of the Tule River have been extensively developed. This development made it possible to disperse much of the heavy flows for preseason irrigation of permanent crops, such as orchards and vineyards. Also, check

dams were installed at several locations on the river to increase the holding capacity of the existing channel. This permitted much of the floodwater to percolate into the ground. As a result of these efforts, only 9 500 dam³ (7,700 ac.-ft.) of excess water flowed to the lake during the February through May period.

Kern River Flows

Kern River water that escaped being used for irrigation or recharged to ground water eventually reached the sedimentation basin at the inlet to the Kern River-California Aqueduct Intertie. The Intertie was constructed by the U. S. Army Corps of Engineers in 1976 to allow Kern River floodflows to enter the California Aqueduct for delivery to other areas, instead of flooding land in the Tulare Basin. At the sedimentation basin, some water was directed south to Buena Vista Lake; some north toward Tulare Lake; and then, on April 6, the Department of Water Resources opened the gates to the Intertie for the first time since its construction in 1976 and let Kern River waters flow into the California Aqueduct.

Once reaching the Aqueduct, the water was routed both southward to Southern California and northward to other service areas. Six pumps were installed at Check 25 in the California Aqueduct north of the Kern River Intertie. The pumps had a capacity of $13.3~{\rm m}^3/{\rm s}$ (470 feet $^3/{\rm s}$) and routed water northward in the Aqueduct to irrigate lands in the Lost Hills Water District and the Buena Vista Water Service District. The pumps were used during the period May 5 to May 26, diverting a total of 2 200 dam³ (18,000 ac.-ft.) of Kern River water away from the Tulare Lakebed. During April, May, and June, a total of 220 000 dam³ (178,000 ac.-ft.) of Kern River water entered the Aqueduct through the Intertie.

Southern California

Between mid-December 1977 and mid-March 1978, an unusually persistent series of storms moved into California from the west and southwest, bringing near-record rainfall and major flows to numerous streams in Southern California. A few streams in San Diego and San Luis Obispo County had their greatest discharge of the season during mid-January, but most peak flows in the area occurred about February 10 and March 4 and 5.

Probably the most serious effects of this series of storms were the numerous landslides, mudslides, and mudflows that resulted from the total saturation of the ground by the extraordinary accumulation of rainfall. In addition to extensive property damage, other damage not evaluated in monetary terms occurred. The deaths of 20 persons can be attributed directly to the flooding, and many other persons suffered hardship and physical injury.

The destruction or damage of numerous hillside and canyon homes --many of them very expensive -- followed the ground saturation and, on several occasions, the mud and landslides blocked major highways, causing long traffic delays. To confound the problems,

communication and power service were frequently disrupted at inopportune times.

Numerous streams throughout the southern portion of the State during and following the torrential downpours encroached or exceeded their carrying capabilities, prompting the evacuation of homes and businesses in low-lying urban and rural areas. Widespread street flooding occurred where drainage systems failed or were not adequate to accommodate the excessive volumes of runoff.

The unusual persistence of the storms made it difficult to relate damage to specific storms. The earlier December and January rains primed the soils and contributed to the increased ground saturation and fast-moving runoff generated by the more intensive storms of February and March. Furthermore, in some cases it was difficult to separate damage caused by floods and that caused by the movement of mud. The total damage in Southern California from February and March floods, not including wind damage, coastal wave action, mudslides or flooding in undeveloped canyon areas and isolated agricultural areas, is estimated at \$86 million.

The flood events of 1977-78 in specific areas of Southern California follow.

Ventura County

The heaviest impact of the January-March storm was in the Santa Clara River Basin along Sespe Creek; however, other areas, including isolated residential and agricultural tracts, suffered substantial material losses and much inconvenience. The total flood damage sustained in Ventura County, primarily during February and March, was estimated to be \$20 million.

Ventura River

The heaviest damage to property along the Ventura River and its tributaries occurred at Ojai, Live Oak Acres, Casitas Springs, and Matilija Dam and vicinity.

At Live Oak on March 4, approximately 26 residential and business buildings were inundated from .30 to 1.5 metres (1 to 5 feet). Some homes, in addition to water damage, lost from 3 to 12 metres (10 to 40 ft.) of property along the river embankment. The effect of the flooding in residential areas was minimized by individual floodfighting efforts, which included sandbagging and elevating home furnishings. Roads, bridges, railroads, and utility structures, however, felt the full impact of the flooding, and the cost of repairing or replacing the facilities was substantial.

In the Matilija Canyon Area, rushing waters damaged parks and ripped out roads and main power lines that served scattered homes in the canyon area. Some homes were damaged, and the severe impact of utility damage was experienced through the loss of service. Another effect of the flooding was the forced closure

of all four lanes of the Main Street Bridge in the City of Ventura. The bridge was not serviceable for two months. The damage to Ventura River drainage areas exceeded \$3.5 million.

Santa Clara River

Flows in the Santa Clara River within Ventura County were generally contained within the channel during the February-March storm. Heavy losses, however, were reported within the flood plain, where dozens of acres of prime agricultural land were washed away, and extensive repairs were required to restore eroded roads, levees, bridges, and utility structures. Unharvested crops were also destroyed as turbulent water washed away the plants or exposed the root system. Prolonged saturation of the soil caused extensive root decay.

Piru Creek

Erosive channel flows during the February and March storms damaged highways and bridge abutments and blocked access to the community of Piru, imposing temporary hardships on some of the residents. The cost of restoring the facilities was estimated at \$1 million.

Sespe Creek

On March 4, severe thunderstorms caused the most damaging flooding on record for Sespe Creek. The evacuation of two residential subdivisions became necessary when Sespe Creek overflowed its banks. Two hundred and four homes and two apartment buildings were inundated to depths of 0.6 to 1.5 metres (2-5 feet) in the Serenos Tract, and eleven business firms suffered substantial losses. Overtopping of the east bank permitted the uncontrolled water to stream through undeveloped land and orange groves; in its course, the floodwater also washed out roads, bridges, and railroad beds, and deposited heavy amounts of silt and debris in industrial and agricultural areas. Access to the city by auto was impossible for several days, necessitating the use of helicopters to move people and supplies during the emergency.

Damage estimates in this flood-stricken area exceeded \$6 million. One life was lost.

Santa Paula Creek

Santa Paula Creek was generally contained within the channels during the high waters of February and March. This is possibly attributable to the channelization and levee work performed in Santa Paula Creek by the Ventura County Flood Control District in early February in anticipation of continued rains and flooding in the Fillmore-Santa Paula areas. The U. S. Army Corps of Engineers conducted floodfight activities during both the February and March storms, and the damage was generally limited to erosion.

Orange County

The February and March storms caused significant damage in the Santa Ana River and a tributary, Santiago Creek. Most damage was attributable to erosive flows in channels, which damaged roads, bridges, utility lines, parks, and the channels themselves. Apartments and a few businesses were damaged on Fullerton Creek, and Caspers Park was heavily eroded by flows through San Juan and Bell Canyon Creeks.

Flood damage for the County totaled \$8.7 million; over half of this amount was damage to flood control works, channels, and streams.

San Diego County

The February and March rains caused flood damage to business and industry within the county. Highways, bridges, and railroads, as well as utilities, residences, public properties, flood control works, and channels also suffered. Flood damage in the county totaled about \$12 million.

San Diego River Drainage Areas

The upper San Diego River Valley and Mission Valley reaches experienced flooding and flood damage.

Mission Valley, a reach extending about 16 kilometres (10 miles) upstream from Mission Bay, experienced recurrent flooding during February and March. All roadway dip crossings in Mission Valley were intermittently flooded and closed to traffic. In spite of precautions taken to barricade and close flooded and washed out roads, two people died trying to cross the river at Mission Center Road.

Fifteen businesses in Mission Valley suffered flood damage, including one car-leasing firm and two new-car vehicle storage lots where parked automobiles were innundated by up to 1.5 metres (5 feet) of water.

San Vicente Creek flooded when San Vicente Dam overtopped and caused damage estimated at \$1.88 million. Peak flows below the dam, as the result of heavy rains on March 4, were estimated to be $57~\text{m}^3/\text{s}$ (2,000 ft. $^3/\text{s}$). Nearly 100 residences, as well as 57~business firms, were damaged.

San Luis Rey River Drainage Areas

Damage from the flooding of the San Luis Rey River and its tributaries occurred to private residences, roads, waterlines, and sewerlines. The industrial park in the flood plain northeast of Oceanside suffered damage estimated at \$2.4 million.

Ostrich Creek, a San Luis Rey River tributary, flooded in March, washing out sewer and waterlines and damaging 29 homes and 15 businesses.

Los Angeles County

The February and March storms caused damage throughout the canyon areas of the San Gabriel Mountains, particularly at Big Tujunga and Little Tujunga Canyons, Pacoima Wash, and extended up to the Santa Clarita Valley. Most of the damage occurring during these storms was from erosive flows in channels, which damaged roads, parks, channels, and flood control works. Flood damage within the County totaled \$34.5 million, including almost \$32 million for the repair of flood control works and channels and streams.

Los Angeles River Drainage Areas

Damage to the Los Angeles River drainage area was \$18.5 million, including more than \$17.8 million in damage to flood control structures, reservoirs, channels, and streams. Some \$17.1 million of this amount was needed to remove debris (much of which consisted of large boulders) from about 60 debris basins and 10 reservoirs.

Oak Street Drain, a downstream tributary to Temescal Wash, flooded part of Corona in January and again on February 10, 1978. Burned-off upstream watershed vegetation from 1977 fires was a contributing factor to the flooding, which caused damage to mobile home parks in the vicinity of 6th Street on February 10. One hundred and twenty-six trailers were involved. Two mobile home parks were evacuated, and one woman died of a heart attack during the evacuation. Further downstream, twelve businesses were flooded and had to remain closed for up to a week. Oak Street Drain floodwaters poured over the top of the Riverside Freeway, which was closed for several hours while crews attempted to repair the damage. Silt and debris damage was widespread. Damage from the Oak Street Drain flooding amounted to about \$1.2 million.

San Bernardino County

The February and March 1978 flooding did not reach the levels experienced in 1969 when the worst floods in the County's history were recorded. However, damage to channels, roads, bridges, railroads, and utilities in the Santa Ana and Mojave River drainage areas was estimated at \$5.2 million.

On Lytle Creek, one home was completely destroyed by floodwaters, and twelve other homes were damaged. Also, water flowed through five homes in Rancho Cucamonga when Cucamonga Creek left its course in the vicinity of 19th Street because debris clogged the channel.

Santa Barbara County

Silt buildup and erosion in channels and streams accounted for the heaviest flood-related damage within the County, especially in the Santa Maria River and South Coastal streams drainage area.

Agricultural damage occurred as streams meandered from their channels at numerous locations and eroded and deposited silt over hundreds of acres of farmland.

In February, half a dozen buildings in the mountain community of Hidden Springs were washed away by a wave that one resident reported as 9 metres (30 feet) high. In addition to exceptional debris damage, many roads, some bridges, railroad appurtenances, water systems, and homes were damaged. One death occurred when a man left his home to check his corral. He was not seen again.

San Gabriel River Drainage Area

As in the Los Angeles River drainage area, the greatest damage, \$13.3 million, was to flood control structures, channels, and streams. Total flood-related damage in the San Gabriel River drainage area was \$13.5 million. Again, and by far the greatest expense, \$13.0 million was for removing debris from reservoirs, debris basins, and spreading basins.

Riverside County

Flood damage totaling about \$9 million occurred in the Santa Ana River, the Santa Margarita River, and the Whitewater River drainage areas.

The main damage along the Santa Ana River from the February and March flooding was to water works, wells, pumps, and rail lines. Also, agricultural lands within the Prado Flood Control Basin were subjected to high and prolonged water levels.

On March 4, while cleanup operations from previous floodings were still in progress, the worst flooding of the season occurred on Temescal Wash in the vicinity of Corona. Using boards and sandbags, more than 60 people struggled to reinforce a levee before conditions became so dangerous the effort had to be abandoned. The rushing waters poured through the embankment, flooding a trailer park and 20 business establishments to depths of up to 1.2 metres (4 feet). The overflow continued downstream to damage roads, wash out railroad tracks, and spill silt over a wide area. Damage from this flooding approached \$2 million.

Costs of repairs to highways, bridges, and railroads were especially high in the Santa Ynez River drainage areas.

San Luis Obispo County

Only minor damage was reported from residential flooding within the county. A number of roads were flooded, and erosion and sedimentation damage occurred, especially along the lower end of Arroyo Grande Creek. However, by far the heaviest damage was to roads and bridges along the Cuyama Rīver where losses exceeded \$10 million.

River Finds A New Way To The Sea



Muddy water and debris — not cars moved through the parking lot at Sar Diego Stadium yesterday, deposited by the rampaging San Diego River



This aerial photo shows a partially submerged to the ocean More than 2½ inches of rain fell during which claimed the lives of a San Diego couple on their



Two unidentified men, admit on a sea that was formerly the Standard Hotel golf course, ciling to a car until rescues arrive. Several motorists who unwitting ly tried to brave the waters found themselves stranded—or swimming




Staff Photos By Tony Doubek and Jerry Rife

Three young girls, pant legs rolled knee-high in vain attempt to keep their clothes dry, survey the dismal scene to the Fashion Valley parking lot from their own private little island. Access to the center was Private Board only.



Survivors at Hidden Springs Tell How Families Perished



AP Loserphoto

Mrs. Ray Filippi stand in what remains of their 75-acre broccoli ranch. Fifty acres of the field were flooded by the normally dry Salinas River, which also carved away a mature eucolyptus orchard. Officials estimate the river has damaged 20,000 acres.

FIGURE 12. NORTHERN CALIFORNIA REFERENCE MAP FOR HYDROGRAPHS, FIGURES 13-16





FIGURE 13. HYDROGRAPHS OF SHASTA LAKE AND SACRAMENTO RIVER







FIGURE 14. HYDROGRAPHS OF LAKE OROVILLE AND FOLSOM LAKE







FIGURE 15. HYDROGRAPHS OF SMITH AND TRINITY RIVERS






FIGURE 16. HYDROGRAPHS OF EEL AND RUSSIAN RIVERS









FIGURE 17. HYDROGRAPHS OF SAN JOAQUIN AND NAPA RIVERS






APPENDIX A

Sacramento River Crest and Weir Overflow Records




A-2, PERIOD OF RECORD OF OVERFLOW OF THE MOULTON WEIR



NOTE:
Onto campiled from records of D.W.R. stream gaging station. Secromenta River at Moultan Weir.
Datum: 0 = 0 U.S.E. D.
Periad of record: 1935 to present
Crest elevation = 76.75 feet (23.41 metres)

Metric Equivalent: 1 FOOT = 0.305 METRES B3 8 feet (25 6 metres)

A-3, PERIOD OF RECORD OF OVERFLOW OF THE COLUSA WEIR



Data compiled from recards of DWR stream gaging station Socramento River at Calusa Weir Datum 0 = 0' U S E D


Period of record: 1935 to present

Crest elevation 61 80 feet (18.85 metres)

Metric Equivalent: | FOOT = 0 305 METRES Designates periods of flow over weir

70 6 feet
(21.5 metres)

A-4, PERIOD OF RECORD OF OVERFLOW OF THE TISDALE WEIR




NOTE:
Date compiled from records at D WR stream gaging station. "Sacromento River at Tisdale Weir
Datum. 0 = 0' U.S.E.O.
Period of record. 1935 to present
Crest elevation = 4545 feet [13.86 metres)

Metric Equivalent: 1 FOOT = 0.305 METRES LEGEND
Designotes periods of flow over weir

53 3 feet
(16 3 metres)

A-5, PERIOD OF RECORD OF OVERFLOW OF THE FREMONT WEIR




NOTE:
Date compiled from records of D.W.R stream gaging, station "Sacramento River at Freemont Weir, West End"
Datum: 0 = 0' U.S.E.O.
Period of record: 1934 to present
Crest elevation = 33.50 feet (10.22 metres)

Metric Equivolent: 1 FOOT = 0 305 METRES LEGEND
Designates periods of flow over weir
for the state of the state

STATE OF CALIFORNIA
THE RESOURCES AGENCY

DEPARTMENT OF WATER RESOURCES


A-6, PERIOD OF RECORD OF OVERFLOW OF THE SACRAMENTO WEIR



NOTE:
Data compiled from records of D.W.R. stream gaging station
Sacramenta Weir Spill to Yolo Bypass, near Sacramento
Oatum: 0=0' U.S.E.O.
Period of record. 1926 to present
Crest elevation = 24.75 feet (7.55 metres)
Elevation of top of gates = 310 feet (9.46 metres)

Metric Equivalent: | FOOT = 0.305 METRES Designates periods of flaw over weir and total number of gates opened

A-7, PERIOD OF RECORD OF INUNDATION OF THE YOLO BYPASS



NOTE:
Date compiled from records of DWR stream gaging station "Total Bypass near Lisban."
Datum: O=U.S.E.D. Datum
Period of Record. 1914 to Present
Assumed overflow of Bypass of stage above 11.5" (3.51 metres)

on the Lisbon gage

Metric Equivalent:

I FOOT = 0.305 METRES

Designotes period of inundation of Bypass

APPENDIX B

Peak Flows and Stages
at
Selected Streams and Stations in California

INTRODUCTION

Appendix B presents data for selected stations on representative streams of the major hydrogrologic basins of California (Figure 2). Historic data are obtained from USGS Surface Water Records, Department of Water Resources' Bulletin 130, and U. S. Department of Commerce, NOAA, National Weather Service, Daily River Stage publications. Current water year data, obtained from USGS and DWR, are preliminary and are subject to revision.

Stations are listed in a downstream direction along the main stream and tributaries. Stations on tributaries are listed between main stream stations in the order in which the tributaries enter the main stream.

LEGEND FOLLOWS TABLES

PEAK FLOW AND STAGES (METRIC UNITS)

	ORAINAGE	PERIOO	SOURCE	. PRI	EVIDUS MAXIM OF RECORD	UM	•	1977-1978 HATER YEA	ir	1
I I	. 30 KM	RECORD	. RECORD	. DATE .	STAGE . IN METRES .	DISCHARGE IN M /S	. DATE .	STAGE . In Metres .	DISCHARGE	
	• • • • • • • • •			H CDASTAL	AREA					•
SMITH	RIVER BASI	N								
SMITH RIVER NEAR CRESCENT CITY	1577	1931-	USGS	12-22-64	14.8	6,460	12=14-77	10.19	2890	
KLAMA	TH RIVER BA	NISIN								
SHASTA RIVER NEAR YREKA	2053	1933-41	USGS	12-22-64	3.9 4.2(A)	608	1-14-78	2.03	60	
SCOTT RIVER NEAR FORT JDNES	1691	1941-	USGS	12-22-64	7,7(AC)	1,550	12-14-77	4.74	376	
KLAMATH RIVER NEAR SEIAD VALLEY	18078	1912=25 1951=	USGS	12-23-64	10.3(A)	4,670	12-15-77	4.50	829	
SALMON RIVER AT SOMESBAR	1945	1911-15 1927-	บรรร์	12-22-64	14.2(A)	3,770	12-14-77	5.53	897	
KLAMATH RIVER AT ORLEANS	21950	1927-	USGS	12-22-64	23,3(AC)	8,690	12-14-77	7.10	3140	
TRINITY RIVER ABOVE COF CREEK NEAR TRINITY CEN		1957-	USGS	12-22-64	3.7 4.1(A)	588	1=14=78	2.54	233	
TRINITY RIVER	1885	1911-	USGS	12-22-55	8,3(AC)	2,030	7-24-78	1.48	21	
NORTH FORK TRINITY RIVER AT MELENA	391	1911-13 1957-	USGS-DWR	12-22-64	8,5(A)	1,010	12-14-77	5.04	225	
TRINITY RIVER	3726	1931-40 1956-	USGS	12-22-55	13,2(4)	4,870	1-16-78	4,66	586	
TRINITY RIVER	7420	1911-14 1916-18 1931-	usgs	12-22-64	12.3(AC)	6,540	1-17-78	10.16	1760	
KLAMATH RIVER NEAR KLAMATH	31339	1910-26 1950-	usgs	12-23-64	16.9(4)	15,800	12-14-77	0.00	9200	
REOWO	OD CREEK 8	ASIN								
REDWOOD CHEEK AT ORICK	720	1911-13 1953-	USGS	12-22-64	7.3(A)	1,430	12-14-77	5.06	600	
LITTL	E RIVER BA	31N								
LITTLE RIVER NEAR TRINICAD	113	1955=	USGS	1-22-72	4.38 4.8(A)	275	12-14-77	3.09	173	
MAO R	IVER BASIN									
MAD RIVER NEAR FOREST GLEN	370	1953-	USGS	12-22-55	7.5(A)	1,110	1=16=78	3.17	231	
MAD RIVER NEAR ARCATA	1256	1910=13 1950=	usgs	12-22-55	9.1	2,200	1-17-78	4.66	620	
EEL F	RIVER BASIN									
EEL RIVER BELOW SCOTT (NEAR POTTER VALLEY	751	1922-	USGS	12-22-64	7.4(A)	1,590	1-16-78	4.41	447	
EEL RIVER AT VAN ARSOAL DAM NEAR POTTER VALLEY	E 903	1909-	USGS	12-22-64	10.3(A)	1,820	1-16-T8	6,21	594	
OUTLET CREEK NEAR LONGVALE	416	1956=	USGS	12-22-64	9,3(A)	2,210	12-14-77	4.61	438	

PEAK FLOW AND STAGES (ENGLISH UNITS)

I I	DRAINAGE	PERIOD	SOURCE	PR'	FVIOUS MAXIM OF RECORD	UM		1977-197	7.8
I STHEAM AND STATION I	. AREA IN . SO MILES	. RECOND	. AECORD	OATF	. STAGE . . IN FEFT .	OTSCHARGE IN CFS	. DATE .	STAGE IN FEET	DISCHARGE IN CFS
							**********	******	
			тепи	H COASTAL	AREA				
	RIVER MAS	IN							
NEAR CRESCENT CITY	609	1931-	usgs	12-22-64	48.5	228,000	12-14-77	33,44	102000
KLAM	TH RIVER R	ASIN							
SHASTA RIVER NEAR YREKA	793	1933+41 1944+	usgs	12-22-64	12.9 13.9(A)	21,500	1-14-78	6.65	2150
SCOTT RIVER NEAR FORT JONES	653	1941+	USGS	12-22-64	25.3(AC)	54,600	12-14-77	15.56	13300
KLAMATH PIVER NEAR SFIAD VALLEY	6980	1912-25 1951-	USGS	12-23-64	33,8(A)	165,000	12-15-77	14.78	29300
SALMUN RIVER AT SUMESHAR	751	1911-15 1927-	usgs	12-22-64	46.6(A)	133,000	12-14-77	18,15	31700
KLAMATH RIVER AT ORLEANS	8475	1927-	uses	12-22-64	76.5(AC)	307,000	12-14-77	23.30	111000
TRINITY PIVER ABOVE CO		1957-	USGS	12-22-64	12.3 13.4(A)	20,800	1-14-78	8.32	8250
TRINITY WIVER	728	1911-	USGS	12-22-55	27.3(AC)	71,600	7-24-78	4.86	750
NORTH FORK TRINITY RIVER AT HELENA	151	1911-13 1957-	USGS-DWR	12-22-64	27,9(4)	35,800	12-14-77	16.52	8000
TRINITY PIVER NEAR HURBI RANCH	1439	1931-40 1956-	USGS	12-22-55	43,2(A)	172,000	1-16-78	15.28	20700
TRINITY WIVEW	2865	1911-14 1916-18 1931-	usas	12-22-64	40.3(AC)	231,000	1-17-78	33.33	65500
KLAMATH PIVER NEAR KLAMATH	15100	1910+26 1950+	USGS	12-23-64	55,3(4)	557,000	12-14-77	0.00	325000
RF (I w	OOD CREEK H	BASIN							
REDWOND CHEEK AT ORICK	278	1911=13 1953=	USGS	12-22-64	24.0(4)	50,500	12-14-77	16,60	21200
LITT	LF PIVFR RA	NS1N							
NEAR TRINIDAD	40	1955+	USGS	1-22-72			12-14-77	10.14	6150
MAD	RIVER HASTI	1							
MAD KIVEH NEAH FOREST GLEN	143	1953•	USGS	12=22=55	24,5(4)	39,200	1-16-78	10.41	8200

12-22-55

12-22-64

12-22-64

12-22-64

29.8

24.2(4)

33.9(4)

30,6(4)

77,800

56,300

64,100

77,900

1-17-78

1-16-78

1-16-78

12-14-77

15,29

14.48

20.38

15.12

21900

15800

21000

15500

1910-13

1950-

1922+

1909-

1956-

FEL HIVER BASIN

290

161

USGS

USGS

USGS

USG5

MAD HIVEH

DUTLET CHEEK NEAR LUNGVALE

BEL RIVER BELOW SCOTT DAM NEAR PUTTER VALLEY

BEL HIVER AT VAN AHSDALE DAM NEAR POTTER VALLEY

PEAK FLOWS AND STAGES (CONTINUED) METRIC UNITS

			METRIC UNIT	5				
I I	. DRAINAGE . PERIDI		. PRE	VIDUS MAXIM OF RECORD	UM	•	1977-1978 WATER YEA	I R I
I STREAM AND STATION I	. SO KM . RECORI	D . RECOR	D. DATE .	STAGE . IN METRES .	DISCHARGE IN M /S	DATE . I	STAGE . N HETRES .	DISCHARGE I IN M /S I
			RTH COASTAL A	REA (CONTIN		•••••		•••••••••
EEL	RIVER BASIN							
EEL RIVER	NTINUED)							
AT FORT SEWARD	5457 1955-	USGS	12-22-64	26,6(AC)	15,900	1-16-78	10.68	3200
SOUTH FORK EEL RIVER NEAR MIRANDA	1390 1939-	USGS	12-22-64	14.0(A)	5,640	1-16-78	6.79	1500
BULL CREEK NEAR WEDTT	72 1960-	USGS	12-22-64	6.3(AC)	184	12-14-77	3,31	120
EEL RIVER AT SCOTIA	8062 1910-	usgs	12-23-64	21.9(A)	21,300	1-17-78	10.61	4790
VAN DUZEN RIVER NEAR BRIOGEVILLE	574 1950-	USGS	12-22-64	7.3(4)	1,360	12-14-77	4.59	529
MATT	OLE RIVER BASIN							
MATTOLE RIVER NEAR PETHOLIA	1911-13 622 1915-	USGS	12-22-55	9,0(C)	2,560	12-14-77	5.87	1180
	HIVER BASIN							
NDYO RIVER NEAR FORT BRAGG	274 1951-	USGS	12-22-64	8.0	679	3 -8-78	4.04	127
	RRO RIVER BASIN							
NAVARRO RIVER NEAR NAVARRO	784 1950-	USGS	12-22-55	12.4(0)	1,630	1-16-78	7.73	637
	IIAN HIVER BASIN							
RUSSIAN RIVER NEAR UKIAH	1911-13 259 1952-	USGS	12=21=55	6.4	535	12-14-77	5,65	280
EAST FORK RUSSIAN RIVE NEAM CALPELLA	238 1941-	USGS	12-22-64	6.2	529	1 -9-78	5,36	230
RUSSIAN RIVER NEAR MOPLAND	937 1939-	USGS	12-22-55 1237	8.2 9.1(A)	1,270	1-16-78	5,25	506
RUSSIAN RIVER NEAR CLOVERDALE	1302 1951=	USGS	12-22-64	9,6(0)	1,560	1=16=78	6.04	784
RUSSIAN RIVER NEAR HEALOSBURG	2053 1939-	USGS	12-23-64 1237	8.2 9.4(A)	2,020	1-16-78	5.88	1180
DRY CREEK NEAM CLOVENDALE	227 1941-	USGS	12-22-64	5.5	512	1=16=78	3,88	272
DRY CREEK NEAR GEYSERVILLE	419 1959-	USGS	1-31-63	5.3	917	1 -9-78	4.88	523
RUSSIAN RIVER NEAR GHERNEVILLE (HACIENDA	A HR.1 3471 1939-	usgs	12-23-64 12-23-55	15.1(A) 15.1(A)	2,640	1=17=78	12.41	1850
		3	N FRANCISCO I	BAY AREA				
-	KER CREEK RASIN							
WALKER CREEK NEAR TOMALES	95 1959-	USGS	1-16-73	7.0	186	1-16-78	6.00	104
COR	TE MADERA CRFEK BASI	IN						
CORTE MADERA CREEK AT ROSS	46 1951-	USGS	12-22-55	5.3	102	1-14-78	4.36	61
NOV	ATO CREEK BASIN							
NOVATO CREEK NEAR NUVATO	46 1946-	USGS	1-14-70	3.4	56	1-16-78	3.55	52

PEAK FERWS AND STAGES (CONTINUED) ENGLISH UNITS

ſ					VIDUS MAXIM			1977-197	
I I STHEAM AND STATION	. DRAINAGE . AREA IN			•	OF RECORD		:	WATER YE	AR
	. SQ MILES	. RECORD	RECORD	. DATE .	STAGE . IN FEFT .	DISCHARGE IN CFS	. DATE .	STAGE . IN FEET .	DISCHARGE IN CFS
			NORT	H COASTAL A	REA (CONTIN	UED)			
	REVER HASIN NTINUED)								
THE THE THE TANKS	2107	1955-	usgs	12-22-64	A7.2(AC)	561,000	1-16-78	34.79	113000
OUTH FORK EEL FIVER NEAR MIRANDA	537	1939-	USGS	12-22-64	46.0(A)	199,000	1-16-78	22.29	52900
POLL CREEK NEAR WEDIT	28	1960-	uses	12=22=64	20.6(AC)	6,520	12-14-77	10.87	4250
EL RIVER AT SCOTIA	3113	1910-	uses	12=23=64	72.0(A)	752,000	1-17-78	35.47	169000
NEAR BRINGEVILLE	555	1950-	USGS	12-22-64	24.0(4)	48,700	12-14-77	15.05	18700
WATT	OLE RIVER HA	81º							
NEAR PETROLIA	240	1911=13 1915=	USGS	12-22-55	29.6(0)	90,400	12=14=77	19.27	38800
WOYD	HIVER BASIS	4							
OYO HIVEH NEAR FORT HRAGG	106	1451=	uses	12-22-64	26.3	24,000	3 -8=78	13.24	4500
NAVA	BHU BIAFE BY	ASI^!							
AVARHO WIVER NEAR WAVARRO	303	1950•	USGS	12-22-55	40.6(0)	64,500	1-16-78	25.35	22500
RUSS	IAN PIVER B	ASIN							
NEAR HKIAH NEAR HKIAH	100	1911=13 1952=	USGS	12-21-55	21.0	18,900	12-14-77	18.54	9900
AST FUME HUSSIAN PIVE NEAR CALPFLLA	65	1941-	USGS	12=22=64	50.5	18,700	1 -9-78	17.57	A150
NEAR HOPLAND	362	1939-	USGS	12=22=55 12= =37	27.0 30.0(A)	45,000	1=16=78	17.23	17900
NEAR CLOVERDALE	503	1951-	USGS	12-22-64	31,6(0)	55,200	1-16-78	19.80	27700
NEAH HEALUSHING OSSIAN HIVER	793	1939-	usgs	12=23=64 12= =37	27.0 30.8(A)	71,300	1=16=78	19.28	41800
WEAR CFUNEHUALE	АН	1941-	uses	12-22-64	18.1	18,100	1-16-78	12.74	9600
NEAR CEASERAILLE	162	1959=	11868	1=31=63	17.5	32,400	1 =9+78	16.00	18500
COEF VEVILLE CHACTENDA	э нь.э 1340	1939-	Pasu	12-23-64	49.6(A) 49.7(A)	93,400	1-17-78	40.73	65200
			SAN	FRANCISCO F	AAY AREA				
« AL »	CER CREEK HA	SIN							
ALKER CHEFK DEAR INMALES	37	1959=	HSGS	1-16-73	22.9	6,600	1-16+78	19.67	3700
(0 + 1	TE MADERA CR	FFK MASIN							
CORTE MADERA CREEK AT RUSS	1.8	1951=	uses	12=22=55	17.5	3,620	1-14-78	14.32	5500
MINA	ATO CREEK RA	51h							
WEAR MINATO	1.6	1946-	USGS	1-14-70	11.0	2,000	1=16=78	10,56	1850

PEAK FLOWS AND STAGES (CONTINUED) METRIC UNITS

	DRAINAGE		SOURCE		TOUS MAXIMUM OF RECORD		. 1977-1978 . WATER YEAR			
I STHEAM AND STATTON I I	. \$0 KM			. DATE .	STAGE . !	IN M /S	DATE . S	TAGE .	DISCHARGE IN M /S	
				FRANCISCO B						
\$0	NOMA CREEK HAS	E T e	200	EMBNC 1200 BY	17 AMEA (CLIN	11000)				
SUNOMA CHEEK	AGA CHEEL HA	, ,								
AT AGIA CALIFNIE	150	1955-	USES	12-22-55	5,2(0)	251	1-14-78	4,16	201	
N: A	PA RIVER HAST	4								
NEAR ST. HELFNA	800	1939-32	USGS	12-22-55	4.9	356	1-16-78	4,93	382	
NEAR NAPA	564	1929-32	USGS	1=31=63	A . 4	478	1-16-78	6.57	433	
PA	CHECO CHEEK H	SIN								
SAN HAMON CREEK AT SAN HAMON	15	1952=	usas	10=13-62	5.2	45	1 -4-78	1.96	16	
SA	N LORENZO CREI	K HASIN								
SAN LORENZO CHEEK AT HAYGARD	98	1939-40	USGS	10=13=62	6.0(A) 6.3(A)	211	1-16-78	3,76	68	
Δ٤	AMEDA CREEK 4.	181h		11-22-77	0,3(-1					
THEDAIL WILL HI										
NEAM PLEASANTIN	365	1962-	USGS	2- 1-63 1-18-73	2.6(C) 3.8	49	1-17-78	3.40	21	
ARROYD VALLE NEAR LIVERMORE	340	1912-30 1957-	USGS	12-23-55	4.2(A)	515	3 -5-78	1.70	30	
ARROYO VALLE AT PLEASANTON	442	1957-	USGS	4- 3-5A	7.7	319	3 -7-78	3,31	25	
ALAMEDA CHEFK NEAM NILES	1639	1891-	USGS	12-23-55	4.5	821	1-16-78	5.53	112	
PATTERSON CHEEK AT INTO CITY		1954-	uses	2- 1-63	6.2(4)	297	1-16-78	3.26	97	
rn	YUTE CHEEK BA	S 1 N								
CHYOTE CHEEK NEAW MAINHUNE	507	1902-12	uses	3- 7-11		707	5-28-78	0.76	2.5	
MPPER PENTIENCIA CHE AT BAN JOSE	F * 56	1961-	USGS	1-21-67	1.9	424	1-14-78	1.70	24	
	IACALIIPE HTVER			,					·	
GUADALUPE PIVER AT SAN JUSE	372	1929-	USGS	4- 2-58	5.1	259	1-14-78	3,37	5.7	
SAMATUGA CEFER	23	1933-	USGS	12-22-55	5.0(C)	77	1-14-78	2.04	73	
42	TANERO CREEK	HAS15								
MATABERS CREEK	1 #	1952-	USGS	2-27-73	1.7	31	2 -7-78	0.83	12	
54	N FRANCISQUIT	lı .								
SAM FRANCISMUTTO CRE AT STANFORD UNIVERS		1950-41	USGS	12-22-55	4.1	157	1-16-78	1.95	66	

PEAK FLOWS AND STAGES (CONTINUED) ENGLISH UNITS

STREAM AND STATE	DRATNAGE	. PERIOD . S	. SOURCE	OF RECORD			. 1977-1978 . WATER YEAR			
SINGUA AND SINII		, SU FILES	. PECOPO	. PECORD	. DATE	. STAGE . . IN FEET .	DISCHARGE IN CFS	. DATE .	STAGE IN FEET	. DISCHARGE . IN CFS
						PAY AREA (CDA				
	SUNUN	CPEEK RAS	i1N							
SUNOMA CREEK										
AT AGIIA CALTENTE		5A PIVER HASIN	1955=	USGS	12=22=55	17.1(C)	8,880	1-14-78	13.66	7100
	40011	4164 - W211								
MEAR ST. MELENA		Ai	1939-32	USGS	12-22-55	16.2	12,600	1=16=78	16.19	13500
NEAR NAPA		218	1929-32	USGS	1=31=63	27.6	16,900	1-16-78	21.54	15300
	PACHE	CO CREEK BA	SIN							
SAN RAMON CREFK AI SAN RAMON		6	1952-	USGS	10=13=62	17.0	1,600	1 -4-78	6,43	600
	SAN LI	DRENZO CREE	r HASIN							
SAN LURENZO CHEEK AT HAYWARD		38	1939-40	USGS	10=13=62	19.7(A) 20.8(A)	7,460	1=16=78	12,32	2400
	AL AME	DA CPEEK PI	SIN		12-22-37	20,11(2)				
ARBOYO MOCHO										
NEAR PLEASANTIIN		141	1962=	USGS	2- 1+63 1-18-73	8.60(C) 12.4	1,760	1-17-78	11.17	950
ARRUYU VALLE NEAR LIVERMORE		147	1912-30 1957-	uses	12=23=55	13.9(A)	18,200	3 =5=78	5.58	1100
ARROYO VALLE AI PLEASANTON		171	1957=	USGS	4= 3=58	25.4	11,300	3 -7-78	10.87	900
NEAR MILES		633	1891=	USGS	12=23=55	14.9	29,000	1-16+78	7.33	3950
AT HAIUN CITY			1958-	USGS	2= 1=63	20.4(A)	10,500	1-16-78	10.71	3450
	COYOT	E CREEK HA	516							
COYDTE CHEEK NEAR MADRONE		196	1902=12	uses	3- 7-11		25,000	5-28-78	2.49	090
IPPEH PEHITEN(14 L	REFK	55	1961-	USGS	1-21-67	6.2	15,000	1-14-78	5,59	850
	GUANA	LUPE WIVER	RASIN							
GHANALHPE WIVEH		144	1929=	11565	4= 2=58	16.6	9,150	1-14-78	11.05	200
SAPATOGA CHEEK AT SARATOGA		Q	1933=	USGS	12=22=55	6.4(C)	2,730	1-14-78	6.69	2600
	~ A T ∆ ()	FRO CREEK	11249							
MATABERD CREEK AT PALO ALTO		7	1952=		2=27=73	5.5	1,100	2 -7-78	2.73	450
		RANCISQUIT	r							
SAF FWANTISHITE (WEEK		1930-41	บริธิร	12-22-55	13.6	5,560	1=16=78	6.40	2350

PEAK FLOWS AND STAGES (CONTINUED) METRIC UNITS

I I I STREAM AND STAT	TON			SOUPEF	•	VIOUS MAXIMO DE RECDED			1977-1978 WATER YEAR		
I I		SU KM			. DATE .	STAGE .	OISCHAPGE . IN M /S .	DATE .		DISCHARGE	
				CENT	PAL CDASTAL	ARFA					
	00 A039	D CREEK BA	SIN								
REDWOOD CREEK AT REDWOOD CITY		5	1959+	USGS	1-31-63	2.9	18	1-16-78	1,77	7.5	
	PESCAD	ERD CREEK I	HASIN								
PESCADERO CREEK NEAR PESCADERO		119	1951=	USGS	12-23-55	6.5	266	1-14-78	3.90	114	
	SAN LO	RENZO RIVE	BASIN								
SAN LURENZO RIVER AT BIG TREES		287	1936=	USGS	12-23-55	6.9	860	1=14=78	6,55	305	
	SOQUEL	CHEEK AAS	IN								
SOGUEL CAFFR AT SOGUEL		104	1951-	USGS	12-23-55	6.A	447	1-16-78	3,59	113	
	PAJARI	HIVER BAS	I M								
BODFISH CREEK NEAR GILROY		18	1959-	USGS	1-31-63	2.5	35	2 -9-78	1.81	10	
THES PINUS CREEK NEAR TRES PINOS		533	1939-	USGS	4+ 4-4]	2,4	228	2-10-78	2.96	176	
SAN RENITO RIVER NEAR HOLLISTER		1517	1949.	USGS	4- 3-58	5.0	328	2 =9=78	4.51	175	
PAJAHU RIVER AT CHITTENDEN		3071	1939-	USGS	12-24-55	9,9	679	2 -9-78	6.55	565	
CORRALITOS CREEK AT FREEDOM		72	1956*	USGS	12-22-55	4.8(4)	105	1-16-78	2.56	37	
	SALINA	S PIVER RA	SIN								
SALINAS HIVER NEAR PO70		181	1942-	USGS	1-25-69	4,2(C) 4,7(A)	526	3 -4-78	6.19	256	
JACK CREEK NEAR TEMPLETON		64	1949-	USGS	2-24-69	3,4	231	2 -8-78	2.49	A7	
ESTPELLA RIVER NEAR ESTRELLA		2387	1954-	USGS	2-24-69	3,2(4)	920	2-10-78	3.18	903	
SALINAS HIVER NEAR HRADLEY		6565	1948-	USGS	2-24-69	6.2(A)	3,310	2-10-78	5.69	2000	
ARROYO SECU NEAR SOLEDAD		631	1901-	USGS	4- 3-58	5.0	801	2 -7-7A	3.82	518	
SALINAS RIVER NEAR SPHECKELS		10763	1929-	usas	2=26=69 1=16=52	8.1(C) A.2(AC)	2,350	2-11-78	6.91	1630	
	CARMEL	RIVER HAS	IK								
CARMEL RIVER AT WINLES DEL MI	n	499	1957-	USGS	4- 2-5A	3.2	201	1-16-78	3.19	199	
	410 00	F RIVER HA	C TA		12-23-55	3.6(4)	196				
HIG SHE RIVER	-10 St	W WINEW HY	01~								
MEAN AIG SUR		151	1950-	USGS	4- 2-58	3,5	160	1 -5-78	4.36	305	

PEAK FLOWS AND STAGES (CONTINUED) ENGLISH UNITS

[DN:	DRAINAGE AREA IN	PERIOD	SDURCE	PRE	VIDUS MAXIMI OF RECORD		• • • • • • • • • • • • • • • • • • • •	1977-197 WATER YE	EAR
		SO MILES	RECORD	RECORD	. DATE .	STAGE . IN FEET .	DISCHARGE . IN CFS .			DISCHARGE IN CES
				CENT	RAL CHASTAL	AREA				
	HEDWOO	D CREEK RA	31N							
REDWOOD CREEK AT REOWDOD CITY		5	1959-	USGS	1=31=63	9.4	644	1-16-78	5.80	250
	PESCAL	ERO CREEK	RASIN							
PESCADERU CREEK NEAR PESCADERO		46	1951-	USGS	12-23-55	21.3	9,420	1-14-78	12.78	4050
	SAN LE	RENZD RIVE	R BASIN							
SAN LORENZO RIVER AT BIG TREES		111	1936-	USGS	12-23-55	9.55	30,400	1 - 1 4 - 7 8	21.48	10800
	SDQUE	. CREE* HAS	IN							
SURULL CHEEK AT SURUEL		40	1951+	USGS	12-23-55	22.3	15,800	1=16=78	11.77	4600
	PAJAR	RIVER HAS	IN							
HODFISH CHEEK NEAR GILRDY		7	1959=	USGS	1-31-63	A.3	1,240	2 -9-78	5,93	350
TRES PINOS CHEEK NEAH TRES PINOS		206	1939-	USGS	4-4-41	7.8	8,060	2-10-78	9.70	6200
SAN BENJIO RIVER NEAR HOLLISTER		586	1949-	USGS	4- 3-58	16.3	11,600	2 -9-78	14.80	6500
PAJARO PIVEP AT CHITTENDEN		1186	1939-	USGS	12+24+55 4+ 3+58	32.5 33.1	24,000	2 -9-78	21.50	8300
CURPALITUS CHEEK AT FREEDUM		28	1956=	USGS	12-22-55	15.6(A)	3,620	1-16-78	A.39	1300
	SALIN	AS PIVER HA	SIF							
SALINAS HIVER NEAR POZU		7 0	1942-	usgs	1-25-69 1-25-69	13,9(C) 15,5(A)	18,600	3 =4=7A	20.30	9050
JACK CREEK NEAR TEMPLETUM		25	1949-	USGS	2-24-69	11.3	8,160	2 -8-78	8.16	3100
ESTRELLA RIVER NEAR ESTRELLA		955	1954-	USGS	2-24-69	10.4(A)	32,500	2=10=7A	10.43	31900
SALINAS PIVER NEAR BRADLEY		2535	1948-	USGS	5-54-69	2n.3(A)	117,000	2-10-78	18.68	70600
APROYU SECU NEAR SULEDAD		244	1901-	USGS	4- 3-58	16.4	28,300	2 -7-78	12.52	18300
SALINAS HIVEH NEAR SPRECKELS		4156	1900-01	USGS	2-26-69 1-16-52	26.5(C) 26.9(AC)	83,100	2-11-78	22.66	57400
	CAHME	L RIVER HAS	1 N							
CARMEL HIVER AT RORLES DEL HIO	7	193	1957=	USGS	4- 2-58 12-23-55	10.5 11.7(A)	7,100	1=16=78	10.47	7050
	HIG S	UR RIVER H	SIN							
RIG SUR RIVER NEAR BIG SUR		47	1950-	USGS	4- 2-58	11.6	5.680	1 -5-78	14.30	10800

PEAK FLOWS AND STAGES (CONTINUED) METRIC UNITS

I STREAM AND STATION . I .		OF . DF ECDRD . RECO		STAGE	010044000			
		• • • • • • • • • • • • • • • • • • • •		IN METRES .	DISCHARGE . IN M /S .			DISCHARGE IN M /S
		c	ENTRAL COASTA	L AREA (CONT	INUED)			
ARROYO	DE LA CRUZ 6	ASIN						
ARROYO DE LA CRUZ NEAR SAN SIMEON	106 19	50= USGS	12- 6-66	4.7	996	1=16=78	3,53	413
ATMA	MARIA RIVER R	ASIN						
SISCHUC HIVER NEAR GARFY	1219 19	40- USGS	1-25-69	4.0	693	2-10-7R	3.60	475
SANTA MARȚĂ RIVER AȚ GUADALUPE	4509 19	40- USGS	1-16-52	2.5(C)	928	3 -4-78	0.00	628 (E
SANTA	YNEZ RIVER BA	91N						
SANTA YNEZ RIVER HELOW GIRHALTAR DAM NEAR SANTA HARBARA	559 19	ene usgs	1-25-69	7.9	1,530	3 =4=78	0.00	481 (F
SANTA CRUZ CREFK MEAR SANTA YNEZ	191 19	41- USGS	2-24-69	4.4(A)	199	2 -9-78	3.77	1 4 3
SAN JO	SE CHEEK HAST	N						
SAN JUSE CHEEK NEAM GOLETA	15 19	41- USGS	1=25=69	3 • 1 3 • 9	56	1=16=78	2,45	50
ATASCA	DERT CREEK RA	81N						
ATASCADENU CREEK NEAR GULETA	49 19	41- USGS	1-25-69	4.0	148	1-16-78	8.58	122
CARPIN	TER1A CREEK H	A 9 1 N						
CARPINTERIA CREEK NEAR CARPINTERIA	33 19	41• USGS	12-27-71	4.3(A)	251			N/
			BOUTH COASTAL	AREA				
	A CPEFK RASIN							
MATTLIJA LHEEK AT MATTLIJA HUT SPHINGS	142 19	27 - USG8	1-25-69	5.0	566	3 -4-78	4.24	467
VENTUHA HIVEH NEAH MEILERS (IAKS	196 19	59- USGS	1-25-69		792(E)	3 -4-78	0.00	562
COYOTE CHEEK NEAH DAM VIEW	33 14	58- USGS	1-25-69	3.7	556	3 -4-78	3.44	173
NENTURA HIVER ARUTURA RABA		1-14 29- USGS	1-52-49	7.4(A)	1,640	2-10-78	5.83	1800
	CLARA HIVER H	ASIN						
SAN CLARA RIVER AT LOS ANGELES-VENTURA CO. LIN	IF 1667 19	52- USGS	1-25-69	5.A	1,950	2 -9-78	3,31	645
PIRU CHEEK ABOVE LAKE PIRU	963 19	55- USGS	2-25-69	5.7(4)	883	3 -4-78	3,11	311
SESPE CREEK NEAR FILLMORE		1-13 27- USGS	1-25-69	6.3 7.6(A)	1,700	2-10-78	6.83	2070
SANTA PAULA CREEK NEAH SANTA PAULA	104 19	27= USGS	2+25=69	4.6(4)	594	2-10-78	4.21	453
WALTHU	CHEEK BASIN							
MALINU CHEEK AT CHATER C NEAM CALAHASAS	271 19		1-25-69	6.5	957	3 -4-7A	5.07	549
	A CREEK RASIN							
HALLUN RALLUNA CHEEK	C-11V UB21V							

				ENGLISH UNITS					
I I SIRLAM AND STATIO	ADEA TH	E PERIOD	SOURCE	· PREV	VIOUS MAXIM			1977-197 WATER YE	AR I
I	. SO MILE	S . RECORD	RECORD	DATE	STAGE .	DISCHARGE .	DATE .	STAGE .	DISCHARGE I
1	•	••••••		•••••••••••	17 FEET .	IN UP5 .		IN FEET .	IN CFS I
			CEN	TRAL COASTAL	AREA (CONT	INUEO)			
A	RROYO DE LA CI	RUZ HASIN							
ARROYH DE LA CHUZ	41	1950-	USGS	13 1-11	15.1	15 200			
NEAR SAN SIMEUN	AI ANIA MARIA RI		0363	12- 6-66	15.3	35,200	1-16-78	11.59	14600
	ANIA PARIA AI	AER JUSTIN							
SISQUEC PIVEP NEAR GAMEY	471	1940-	USGS	1-25-69	13.0	24,500	2+10-78	11.80	16800
SANTA MARIA RIVER AT GUAPALUPE	1741	1940-	usgs	1-16-52	8.2(0)	32,800	3 -4-78	0.00	22200(E)
S	SANTA YMEZ HIV	ER BASIN							
SANTA VINEZ RIVEP RELIJW GIMPALTAN DA NEAR SANIA RANHARA	M 216	1920-	USGS	1=25=69	25.8	54,200	3 -4-78	0.00	17000(E)
SANTA CRUZ CREEK NEAR SANTA YMEZ	74	1941-	USGS	2-24-69	14.5(4)	7,050	2 -9-78	12.37	5050
5	SAN JUSE CREEK	HASTN							
SAP JUSE CHEFK NEAH GOLETA	Ь	1941-	usas	1-25-69	10.1 12.7	2,000	1-16-78	8.05	1750
, and the second second	TASCADEPO CRE	EK RASIN							
ATASCADERO CREEK NEAR GOLETA	19	1941-	USGS	1-25-69	13.0	5,230	1-16-78	8.47	4300
(ARPINIFHIA CF	FE+ HASIN							
CAPPINIERIA CHEEK	1.5	1441-	1151-5	12-27-71	14.1(A)	8,880			N/A
			sou	TH COASTAL A	kFΔ				
	ENTINA CREEK	HASIN							
MATICIJA CHEEK AI MATICIJA HIII SE	PRINGS 55	1927-	usgs	1-25-69	16.5	20,000	3 -4-78	13,91	16500
NEAR MEINERS (ARS	76	1959-	USGS	1-25-69		28,000(E)	3 -4-78	0.00	19900
CUADIF CHEFK	13	1958-	USGS	1=25=69	12.0	8,000	3 -4-78	11.28	6150
VENTURA FIVER	1 48	1911-14	uses	1=25=69	24,3(A)	58,000	2-10-78	19.14	63600
:	SANIA FLAFA PI	VER PASIA							
SAN CLAPA HIVER AT ANGELES-VENTURA CL		1952-	USGS	1-25-69	34.0	68,800	2 -9-78	10.85	22800
PIRU CREEK ANOVE LAKE PIEU	372	1955=	uses	2-25-69	18,6(A)	31,200	3 -4-78	10.20	11000
SESPE CHEEK		1911-13							
MEAN FILL WHE	251	1927-	U5G5	1-25-69 2-25-69	20.8 25.0(A)	£0,000	2-10-78	22.40	73000
SANTA PALILA CHFEK NEAH SANTA FALILA	40	1927=	USIS	2-25-69	15,2(4)	21,000	2-10-78	13.80	16000
	ALIBU CREEK B	ASI							
MALIMI CHEFK AT CHI NEAM CALAMASAS	ATER CAMP	1931-	IISGS	1=25=69	21.4	33,800	3 -4-78	16.64	19400
	HALLINA CREEK	RASIF							
HALLING CHEEK	90	1428+	USGS	11-21-67	14.9	32,500	2-10-78	14.84	28100

PEAK FLOWS AND STAGES (CONTINUED)

1 .	DRAINAGE	PERIOD	. SOURCE		10US MAXIMU OF RECORD			1977-1978 hater year	,
I STHEAM AND STATION . I . I	SI) KM	. RECORO	. PECDAD	DATE	STAGE . N METRES .	DISCHARGE . IN M /S ,	DATE . :	STAGE , N METRES .	DISCHARGE IN M /S
			รบก ร	H COASTAL AR					
EUS ANE	ELES PIVF	R BASIN							
US ANGELES PIVER AT SEPULVEDA DAM	409	1929-	USGS	1-25-69	3.5	390	3 -4-78	3.67	416
OS ANGELES RIVER AT LOS ANGELES	1331	1929+	USGS	3- 2-38		1,900	2-10-78	4.46	1490
IO HONDU NEAR DOWNEY	370	1928-	USGS	1=25=69	4.6	1,330	3 -1-78	3.73	906
SANTA	NA RIVER	HASTN							
ANTA ANA RIVEH NEAR MENIONE	541	1896-	USGS	3- 2-3R	4.4(0)	1,480	2-10-78	0.00	61
AN GARRIEL RIVER Relow Sapta ee dam Near haedwin park	611	1942•	USGS	1-26-69	6.8	874	3 -5-78	5.55	402
ANTA ANA PIVEH AT "E" SI NEAR SAN HERNARDINN	1377	1939-54 1966-	USGS	2=25=69	5.0	792	3 -4-78	1.74	387
ILL CHEEK NEAR YUCAIPA	108	1919-38	USGS	1-25-69	5.1(A)	1,000	2-10-78	3,37	152
YTEF CHEFK NEAR FUITANA	119	1918-	USGS	1-25-69	4.6(4)	1,020	3 -4-78	2.38	0.0
ANTA ANA HIVEN AT M.W.D. CROSSING	2211	1970-	usas	12-29-70	3,3	150			
AN JACINTO HIVEN NEAR SAN JACINTO	365	1920-	USGS	2-16-27		1,270	1-15-78	4.34	127
ANTIAGO CREEK At MODJESKA	33	1961-	USGS	2=25=69	1,9	184	3 -4-78	0.00	84
ANTIAGO LHEEN Al Santa ama	246	1928-	USGS	2=25+69 1=16=52	2.8(C) 3.0	186	3 -4-78	2.07	52
SAN JU	N CHEEK 8	ASIN							
AN JUAN CEFEK UEAR SAC JUAN CAPISTRANI	274	1928-	USGS	2-25-69	1.7(AC)	634	3 -4-78	0.00	212
	HASIF:								
ANTA MAPLARITA PIVER NEAR TEMECULA	1522	1923=	USGS	2-16-27	4.5(C)	707	3 -1-78	0.00	396
ANTA MANGANITA HIVER AT YSJUDHA	1914	1423-	USGS	2-16-27	5.5(0)	951	3 -1-78	5.52	600
SAN LU	IS PEY RIV	FR HASIN							
AN EUIS HEY HIVER AT Monserate narrows or pag	A GAN	1935-41 1946-	USGS	2- 7-37	2.7(0)	- •	3 -1-78	1.66	47
AN IUIS HEY HIVER NEAF HUNSALL	1326	1916-18	USGS	3- 3-3A	4.9	512	1-17-78	4.56	233
SAN OI	EGUITO PI	FF MASIN							
ANTA YSAMPL CHEFK NEAR PANTIA	290	1912-23	uses	1-27-16	4,3(0)	804	3 -1-78	2.83	113
IANTA YSAHFL CHEFK NEAH SAN PASILIAL	331	1945-12	uses	3-24-06	1.9(0)	226	3 -1-78	3.15	124
SAN DI	een Plate	HASIN							
AN DIEGO MIVER PIEAK SANTEE	976	1412-	USGS	1-27-16	7.7(0)	1,990	1=15=78	3.29	*0
	ATE HIVE								
WEETEATER RIVER NEAF DESCARSO TIJUAN	119 A RIVER BI	1965-27 1956-	uses	2-14-27	4.0(AC)	317	3 =2=7R	2.47	32
IJIIAFA PIVER				3					
NEAR DULYURA	1245	1936=	USGS	2= 7=37 6.2	2.6	133	3 -1-78	2.44	84

		AINAGE . PERIDO	. SOURCE .	. PRE	OF RECORD	ум .	1977-1978 I WATER YEAR J			
I STREAM AND STATION . I .	AREA IN SQ MILES	. RECORD	HECORD	DATE	STAGE . IN FEET .	DISCHARGE .	DATE .	STAGE IN FEET	. DISCHARGE . IN CFS	
			SOUT		REA (CONTIN					
LOS AND	SELES RIVE	R FASIN								
US ANGELES RIVER AT SEPULVEDA DAM	158	1929-	USGS	1=25=69	11.4	13,800	3 -4-7A	12.04	14700	
US ANGELES PIVER AT LUS ANGELES	514	1979-	USGS	3= 2=38		67,000	2-10-78	14.62	52700	
ID MONDU NEAR DOWNEY	143	1928-	usas	1-25-69	15.2	46,900	3 -1-78	12.24	32000	
SANTA	ANA RIVER	RASIN								
ANTA ANA RIVEH NEAR MENTONE	209	1896=	USGS	3- 2-38	14.3(0)	52,300	2-10-78	0.00	2150	
AN GARRIEL PIVER BELUW SANTA FE DAM NEAP BALDWIN PAPK	236	1942-	USGS	1-26-69	55.5	30,900	3 =5=78	18.22	14200	
ANTA ANA HIVER AT °F° S' NEAR SAN BERNARDINO	532	1939=54 1966=	USGS	2=25=69	16.5	28,000	3 -4-78	5.70	13700	
ILL CREEK NEAR YUCAIPA	42	1919-38	USGS	1-25-69	16.8(4)	35,400	2-10-78	11.05	5400	
YTLE CREFK NEAR FONTAMA	46	1918-	USGS	1-25-69	15.0(A)	35,900	3 +4=78	7.80		
ANTA ANA RIVER AI M.W.D. CROSSING	854	1970-	usgs	12-29-70	10.9	5,300			N/A	
AN JACINTH RIVER NEAR SAN JACINIO	141	1920-	usgs	2-16-27		45,000	1=15=78	14.25	4500	
ANTIAGO CHEEK AI MODJESKA	13	1961-	USGS	2-25-69	6.2	6,520	3 -4-78	0.00	3000(F)	
ANTIAGO CREFK AT SANTA ANA	95	1928=	usgs	2-25-69	9.1(C) 9.8	6,600	3 -4-78	6.80	1850	
SAN JU	AN CHEEK F	RASIN								
AN JUAN CREEK NEAR SAN JUAN CAPISTRAM	n 106	1928-	USGS	2-25-69	5.6(AC)	22,400	3 -4-78	0.00	7500	
SANTA : RTVEH	MARGARITA HASIN									
ANTA MARGAMITA RIVER NEAR TEMPCULA	588	1923-	USGS	2=16-27	14.6(0)	25,000	3 -1-78	0.00	14000(E)	
ANTA MARGARITA RIVER AT YSIDURA	739	1923-	USGS	2-16-27	18.0(C)	33,600	3 -1-78	18.12	21200	
SAN LU	IS REY PIV	VER BASIN								
AN LUIS HEY HIVER AT MONSERATE NAHRONS NR PAI	A 373	1935-41	USGS	2- 7-37	A,7(C)		3 -1-78	5.45	1700	
AN LUIS REY RIVER NEAR BUNSALL	512	1916=18	USGS	3= 3=38	16.0	18,100	1-17-78	14.97	4250	
SAN DI	EGUITO RIV	VER HASIN								
ANTA YSAHEL EREEK NEAR RAMUNA	112	1912-23	USGS	1-27-16	14.0(0)	28,400	3 -1-78	9.28	4000	
ANTA YSAHFL CHEEK NEAH SAU PASQUAL	128	1905-12	USGS	3-24-06	6,3(0)	8,000	3 -1-78	10.35	4400	
SAN OI	FUN RIVER	MASIN								
AN DIEGO HTVER NEAR SANTEE	377	1912-	usgs	1-27-16	25.1(0)	70,200	1=15=78	10.79	2850	
	ATER RIVER									
WEETPATER RIVER NEAR DESCANSO	46	1905=27 1956=	usgs	2-16-27	13.2(40)	11,200	3 -2-78	8.12	1150	
	N PIVER H	NS1N								
TIJUANA HIVER NEAR OULZURA	481	1936=	uscs	2- 7-37	A.5	4,700	3 -1-78	8.00	3000	
				63						

				LEINIC UNI	15				
1 1 1 STHEAM AND STATION	. DPATNAGE	PERIOD	. SOURCE		EVIDUS MAXIM OF RECORD		•••••	1977-1978 WATER YEA	9
1	. 50 kM		RECORD	. DATE .	STAGE .	DISCHARGE . IN H /S .	DATE .	STAGE IN METRES .	DISCHARGE
			CENT	AL VALLEY	AREA				
SACHA	MENTIL HIVER	MIBAA							
SACRAMENTI) RIVER AT DELTA	1100	1944•	USGS	12-22-64	6,1	1,100	1-16-78	5.03	874
MEAN HIEHEN	6410	1904-31 1951-	usas	3-19-07	5.1	957	4-29-7A	1,82	54
PIT PIVER RELOW PIT NO.4 FAM	12035	1922=	usgs	1-25-70	5,5	920(E)	3 -9-78	2.92	147
MCCLUID HIVER	1564	1945-	USGS	12-22-55	8.6	1,280	1-16-78	6.79	59
SACRAMENTO RIVER AT KESWILK	16752	1938-	USGS-DWR	2-23-40	14,4(C)	5,270	3 -9-78	7,64	1130
CLEAR CREEK AT FRENCH GULCH	291	1950-	USGS	12-22-64	4.2	215	1-14-78	4.07	258
CLFAH CHEEP NEAR IGU	590	1940-	USGS	12-21-55	4 . 2	693	1-17-78	2,50	160
NEAR WILLAILLE	1100	1949-	USGS	12-27-51	6.6	1,280	1 -9-78	5,58	1020
NEAR CHITCHWOLD	2387	1940-	USGS	12-22-64	6,0	1,700	1 -9-78	5,46	1110
MATTLE CMEEK HELDW COLEMAN FISH HATCHERY NEAR COTTURNOOD	927	1961-	usas	12-11-37	4.8(AC)	991	1-16-78	2,30	167
SACRAMENTO HIVER AT HEND HRIDGE		1960-	Dwk	1-24-70	14.7	4,470	1 -9-78	11.97	2840
PAYNES CHEEK NEAM RED HEURE	240	1949-	USGS	12- 1-61	3.4	300	1-16-78	2.54	122
NED HANK CHEFF NEAR RED HEDEF	243	1948-	DwR	1= 5=65	3.1	275	1-14-78	3.09	263
ANTELUPE CHEEK NEAR HED HLUFF	318	1940-	USGS	1-23-70	5.5	487	1-16-7R	3.87	139
FLORM CHEEK NEAM PASHENTA	240	1948-	USGS	2-24-5A	4,2(0)	331	1-14-78	2.97	182
MEAN LOS MOLINOS	330	1989-13	U568	12-11-37	7.1(4)	1,030	3 -4-7A	3.04	1 89
THOMES CHEEK	512	1920-	USGS=DwR	12-22-44	4.7	1,070	1-14-78	2.29	210
DEER CHEFF HEAR VIIIA	534	1911-15	USGS-DWR	12-10-37	5.9(A)	673	1-16-7A	3,23	236
SACRAMENTO PIVER AT VIVA HEIDGE		1945-	DWR	1-24-70	58.4(1)	4,840 6,460(L)	1+16+78	26.57	3430
SACRAMENTO HIVEH MARILTON CITY (HEFORE SHASIA DAM)		1927-43	DWH	12-11-37	45,9(C7)	4,840(EL)			
SALWAMENTI: WIVEW AT HAWILTING CITY (AFTEN SHASTA LAM)		1944-	DwR	1-24-70	46.0(1)	4,420	1-17-78	14.38	3480
MIG CHICD CMEEK	186	1930-	11565	1= 5=65	4.7	271	1-16-78	3.12	133
STUNY CHEEN NEAR FRUIT	1548	1901-12	USGS	12=23=64	4.8	1,140	1-14-78	3,66	685
SACHAMENTH HIVEH AT HED FEHRY (HEFHRE SHASIA DAM)		1921-43	Dwk	2-28-40	37.1(1)	1,140(EL)			
SACHAMENTII HIVEH AT IJHD FEHRY (AETEM SMASTA IJAM)		1944-	Dwa	1-24-70	36.5(T)	7,500(EL)	1-17-78	20.18	3450

- + 1921-43 USGS-OWF 2- 7-42 29,5 7,500

SACRAMENTO HIVER AT HUITF (ITY (HEPOME SHASTA DAM)

I I I STREAM AND STATION	. DRAINAGE . AREA IN				OF RECORD			1977-19	
I SINCEM AND SIEITON	. SQ MILES		RECORD	DATE .	STAGE .	DISCHARGE .	DATE .	STAGE IN FEET	. DISCHARGE I
			CENT	RAL VALLEY					
SACE	RAMENTO RIVER	RASIN							
SACRAMENTO RIVER AT DELTA	425	1944-	USGS	12-22-64	20.1	38,800	1-16-78	16.51	30900
PIT RIVER NEAR HIEBER	2475	1904-31 1951-	USGS	3-19-07	16.7	33,800	4-29-78	5.98	1900
PIT RIVER BELDW PIT NO.4 DAM	4647	1922-	USGS	1-25-70	18.1	32,500(E)	3 -9-78	9.58	5200
MCCLOUD RIVER ARDVE SHASTA LAKE	604	1945-	USGS	12-22-55	28.2	45,200	1-16-7A	22.27	2100
SACRAMENTO RIVER AT KESWICK	6468	1938+	USGS=DWR	2-23-40	47.2(C)	186,000	3 -9-78	25.05	39800
CLEAR CREEK AT FRENCH GULCH	115	1950-	USGS	12-22-64	13.7	7,600	1-14-78	13.35	9150
CLEAR CREEK NEAR 1GD	228	1940-	USGS	12-21-55	13.8	24,500	1-17-78	8,20	5650
CDW CREEK NEAR MILLVILLE	425	1949-	USGS	12-27-51	21.6	45,200	1 -9-78	18.31	36000
COTTONWOOD CREEK NEAR COTTONWOOD	922	1940-	USGS	12-22-64	19.6	60,000	1 -9-78	17.92	39100
RATTLE CREEK RELOW COLEMAN FISH HATCHERY NEAH COTTONWOOD	358	1961=	USGS	12-11-37	15.8(AC)	35,000	1=16=78	7.54	5900
SACRAMENTO PIVEN AT HEND HHIDGE		1960-	OwR	1-24-70	48.3	158,000	1 -9-78	39.27	100200
PAYNES CREEK NEAR RED BLUFF	93	1949-	USGS	12= 1=61	11.3	10,600	1-16-78	8.33	4300
RED HANK CHEFF	94	1948-	DWR	1= 5=65	10.1	9,730	1-14-78	10.15	9300
ANTELUPE CHEEK NEAR RED HLUFF	123	1940-	USGS	1-23-70	18.0	17,200	1=16=78	12.69	4900
ELDER CREEK NEAR PASKENTA	93	1948-	USGS	2-24-58	13,9(0)	11,700	1-14-78	9.74	6450
MILL CHEEK NEAM LUS MOLINUS	131	1909=13 1928=	USAS	12-11-37	23,4(4)	36,400	3 -4-78	9.97	6700
THOMES CREEK AT MASKENTA	194	1920=	USGS=OWR	12-22-64	15.3	37,800	1-14-78	7.51	7400
NEER CHEEK NEAR VINA	208	1911=15 1920=	บร.เระกพล	12-10-37	19,2(A)	23,800	1=16=78	10.60	A350
SACHAMENTO RIVER AT VINA HRIDGE		1945=	Dws	1-24-70	191,5(7)	171,000 228,000(L)	1-16-78	A7.1A	121000
SACRAMENTO HIVER AT HAMILTON CITY (REFORE SHASTA DAM)		1927=43	DwR	12-11-37	150.7(01	350,000(EL			
SACRAMENTO RIVER AT HAMILION CITY (AFTER SHASTA DAM)		1944-	Dak	1-24-70	150.8(1)	156,000	1 - 1 7 - 7 8	47.18	123000
PIG CHICO CREEK NEAR CHICO	72	1930=	USGS	1= 5=65	15.4	9,580	.1=16=78	10.25	4700
STONY CREEK NEAR FROTH	598	1901=12	USGS	12-23-64	15.9	40,200	1=14=78	12.72	24200
SACHAMENTO HIVEH AT OHO FERRY (HEFORE SHASTA DAY)		1921-43	DNH	2=28=40	121,7(1)	370,000(EL			
SACRAMENTO HIVER AT ORD FERRY (AFTER SMASTA DAM)		1944-	DWH	1-24-70	119.8(T)	265,000(FL	1-17-7A	66,21	122000

96.9 170,000

1921-43 USGS-DVR 2- 7-42

SACRAMENTO RIVER AT BUTTE LITY (REFURE SHASTA DAV)

				-C1416. 041					
1	RAINAGE	PERIOD OF	. SOURCE	PR	EVINUS PAXIMO OF RECORD	. м		1977-197	A A R
1	SO KM		. RECORD .	DATE .	STAGE . IN METRES .	DISCHARGE .	DATE .	STAGE IN METRES	DISCHARGE
********************				RAL VALLEY	AREA (CONTIN	UED)	••••••		
SACRAMEN (CONTIN	TO RIVER UED)	HASIN							
SACRAMENTO HIVER AT HUTTE CITY (AFTER SHASTA DAM)		1944-	USGS-DWR	2+20+58 1+24+70	29.5	4,530 6,730(L)	1=17=78	28.30	3430
MOULION HEIR SPILL TO MUTTE MASIN		1935=	DWR	1=25=70	25,5 25,5	1,030(8)	1-17-78	24,82	521
COLUSA WEIR SPILL TO BUTTE HASIN		1935*	DWR	3- 1-40	21,5	2,440(8)	1=17=78	20.52	1590
SACRAMENTO HIVEH	31365	1940-	USG8=0WR	5 × × × × ×	21.1	1,390	1-18-78	20.23	1280
CULUSA HASIN DHAIN AT HIGHHAY 20		1924-	OMR	2=21=58	15.8	719(E)	1-18-78	15,79	294
HUTTE CHEEK NEAR CHICO	380	1930-	usgs	12=22=64	4.3	600	3 -4-78	2,61	555
HUTTE SLINGS		1968-	D in R	1-26-70	18.7(E)	4.300(E)	1-18-78	17.78	2860
TISDALE WEIR SPILL TO SUITER HYPASS		1940-	DINR	3- 1-40	16.2	727(8)	1-18-78	15.09	47 A
SACRAMENTO HIVER HELUW WILKINS SLODIGH	33478	1938.	USGS	1-26-70	15.5 16.1	A29	1=18=78	14.98	POR
SACRAMENTO HIVER AT KNIGHIS LANDING	37661	1921-39	USGS=DWR	1-26-70 2- 8-42	12.5 12.7(N)	A72	2 =8=78	0.00	841
MIDDLE FURK FEATHER	1776	1925-	USGS	2= 1=63	4.9	410	1=18=78	3.18	9.2
MIDDLE FUHK FEATHER RIVEN HEAR MENHIMAC	2750	1951=	USGS	12-22-64	A.1(A)	2,440	1-17-78	3.82	331
NORTH FORK FEATHER RIVER NEAR PRAITVILLE	1276	1905-	USGS	3-19-07	4,9(0)	283	3-16-78	1.76	3 A
BUTT CREEK HELOW ALMANDM-MOTT CREEK TUNNEL NEAM PRATTVILLE	178	1936=59	USGS	12-23-64	1.4	108	1-14-78	0.59	12
INDIAN CHEEK NEAH CHESCENT HILLS	1914	1906-18 1930-	USGS	3=19=07	6.2(C)	707	3 -5-78	2.78	137
SPANISH CHEEK ABOVE HEACKHANK CHEEK AT KEDDIF	476	1933=	USGS	12-22-64	4 + 1	436	1-16-78	2,52	149
NORTH FURE FEATHER RIVER AT RULGA	505A	1910-	USG5	12-22-64	10.9	2,070(H)	1-15-78	5.91	506
WEST HEAVEH FEATHER RIVER NEAR PAMADISE	245	1457=	USGS=UWR	12-22-64	8.0[4]	744	1-14-78	3,89	171
FEATHER HIVER AT UROVILLE (MEFORE OROVILLE DAM)		1894=67	USG5=DWH	3-19-07	8.6	6,510(CP) 7,140(9)			
FRATHER HRUVILLE DAM) AT CHOVILLE (AFTER HRUVILLE DAM)	9386	1967-	USGS=DWR	1-25-70	и.7	1,590(N)	10 -4-77	0,00	26
HERMALITO AFTERMAY HELFASE TO FEATMER HIVEN MEAN OROVILLE		1967=	USGS-DWR	1-28-10	T . 1	611	3 =9=7A	2,55	454
FEATHER FIVER NEAR GRIDLEY (HEFORE OKOVILLE DAM)	9521	1929-67	USGS-DHR	12-23-55	31,2(1)				
FEATHER WIVEN (AFTER UNIV)(LF DAM)	9520	1967=	IJSGS=DWR	1 = 27 = 70	28,3(1)	2,060	3 =6=7A	24.77	492

5.9

12-26-64

80 1950-

USGS

498

1-16-78

2.73

93

SOUTH MONEHT LMEEK

I I I STHEAM AND STATION	1977-197A I WATER YEAR I			
			STAGE DISCHARGE DAT	TE . STAGE . DISCHARGE I

CENTRAL VALLEY AREA (CONTINUED)										
SACRAMENT (COMTINU		RASIN								
SACRAMENTO RIVER AT MUTTE CITY (AFTER SHASIA DAM)	••	1944-	USGS+DWR	2-20-58 1-24-70	96.7	160,000 225,000(L)	1-17-78	92.84	121000	٠
MOULTON WEIR SPILL TO MUTTE BASIN		1935-	DWR	1-25-70	83.6 83.8	36,400(8)	1-17-78	81.44	18400	
COLUSA WEIR SPILL TO HUTTE BASIN		1935*	DwR	3- 1-40	70,6	86,000(8)	1-17-78	67.32	56100	
SACRAMENTO RIVER AT COLUSA	12110	1940-	USGS-DWR	2- 8-42	69.2	49,000	1-18-78	66.36	45200	
COLUSA MASIN DRAIN AT HIGHPAY 20	•-	1924-	DWR	2-21-58	51.9	25,400(E)	1-18-78	51.81	10400	
NEAR CHICO	1 4 7	1930-	USGS	12-22-64	14.1	21,200	3 -4-78	8.56	7850	
MEAN MERIUIAN	••	1968=	DER	1-26-70	61.5(E)	152,000(E)	1-18-78	58.32	101000	
TISDALE WEIR SPILL TO SUTTEN HYPASS		1940-	0 % R	3- 1-40	53.3	25,700(8)	1-18-78	49.52	16900	
SACHAMENTO HIVEH HELDW WILKINS SCHIGH	12926	1938+	uses	1-26-70 3- 1-40	50.7 52.8	29,300	1-18-78	49.15	28600	
SACRAMENTO PIVER AT KNIGHTS LANDING	14541	1921-39	USGS-DWR	1-26-70	40.9 41.8(D)	30,800	2 -8-78	0.00	29700	
MIDDLE FORK FEATHER RIVER NEAR CLID	686	1925=	USGS	2- 1-63	16.2	14,500	1-18-78	10.44	3250	
MINDLE FORK FEATHER MINDLE FORK MERRIMAC	1062	1951-	USGS	12-22-64	26.5(A)	86,200	1=17=78	12,52	11700	
NORTH FORK FEATHER RIVER NEAR PRATTVILLE	493	19115+	USGS	3-19-07	16,2(0)	10,000	3-16-78	5.77	1350	
BUTT CREEK HELUK ALMADON-HUITT CHEEK TUNNEL NEAK PRATTVILLE	69	1984=	U\$65	12-23-64	5.9	3,830	1 = 1 4 = 7 A	1.94	450	
INDIAN CHEEF NEAR CHESCENT FILLS	739	1906-18	usas	3-19-07	50.5(0)	25,000	3 -5-7A	9.12	4850	
SPANISH CREEK AROVE BLACKHAZIK CREEK AT KEDDIE	184	1933-	USGS	12-22-64	13.5	15,400	1-16-78	8.26	5300	
NORTH FORK FEATHER PIVEN AT PULGA	1953	1910-	USGS	12-22-64	35.A	73,000(H)	1-15-78	19,38	17900	
WEST BRANCH FRATHER RIVER NEAR PARADISE	110	1957=	USGS=DWR	12-22-64	26.2(A)	26,300	1-14-78	12.77	6050	
EFATHER OFFICE DAMS OFFICER OFFICER O	1624	1894-67	USGS=DWR	3-19-07 12-22-64	2.85	230,000(CP 252,000(Q)				
FEATHER FIVEH AT OROVILLE (AFTER OROVILLE DAM)	3624	1967=	US65-0+8	1-25-70	15.3	56,300(N)	10 -4-77	0.00	950	
THERMALITO AFTERMAY RELEASE TO FEATHER RIVER NEAR DEDVILLE		1967=	USGS=0#F	1-28-70	23,3	21,600	3 =9=78	A.35	16000	
FEATHER HIVER HEAR GHIDLEY (BEFORE DROVILLE DAM)	3676	1929-67	IISI,S=PWR	17-23-55	102.2(1)					
FEATHER GROVILLE DAY)	3676	1967=	USGS-DWR	1-27-70	92,8(1)	72,900	3 -6-78	81.25	17400	

12-26-64 19.3 17,600 1-16-78 8.96

3300

SDUTH HURLIT CHEFK
NEAR HARGUR 31 1950= USGS

1			•		PREVIOUS MAXIMUM		1977-1978
1	. DHAINAGE	. PERIOD	. SOURCE	•	OF RECORD		WATER YEAR I
I STREAM AND STATION	. AREA IN	. UF	. UF	,		• • • • • • • • •	
I	. 90 KM	. WECORD	. RECORD	. DATE	. STAGE . DISCHARGE	. DATE	. STAGE . DISCHARGE I
1					. IN METGES . IN M /S		. IN METRES . IN M /8 I

CENTHAL VALLEY AREA (CUNTINHED)

SACRAM	ENTU	RIVER	RASIN

SACRAMENT (CONTINU		PASIN							
FEATMEN NIVER AT YUBA (11Y	10292	1943-	USGS-DWR	12-23-64	23.3 25.1	4,870	3 -7-78	15.21	0.0
NORTH YUBA HIVER BELOW GOUDYFARS HAR	647	1930-	USGS	2- 1-63	7.3(A)	1,130	1-16-78	2.74	129
NOHTH YUHA RIVER BELOW NEW HULLARDS HAR DAM NEAR SAN JUAN	1269	1940-	BSGS	1-22-70	10.8 12.3(C)	1,590 2,590(M)	12-15-77	1.82	0.7
SDUTH YUMA RIVER NEAH CISCO	134	1942=	U\$G\$	1-31-63	6.3(A)	521	5-14-78	2.23	58
SOUTH YUNG RIVEH AT JONES HAR NEAH GRASS VALLEY	797	1940-48	USGS	12-22-64	7.6(4)	1,520	1 -5-78	3.87	233
YUBA HIVER HELDW ENGLEMMIGHT DAM	2869	1941-	U 3 G 3	12-22-64	171.9(0)	4,840(K)	1 -5-78	4.78	45A
DEER LHIEK NEAM SMAHIVILLE	220	1935-	USGS	10-13-62	4.2	328	1 -5-78	2.62	105
YURA HIVEH NEAH MARYSVILLE	3467	1940-	USGS	12-22-64	27.5	5,100	1=16=78	21.28	529
NEAM MMERIFALID	756	1929=	U\$63	12-22-55	5.9(C) 6.3(C)	934	1-17-78	4.00	255
FEATHER HIVEH AT NICULAUS	15333	1943-	USGS-OWH	12-23-55	15.7	10,100	1 =6=78	10.18	741
FREMUNT WELF (MEST END) SPILL TO YOUR HYPASS		1934-	DWH	12-23-55	12.1	A,330(A)	1-18-78	11.10	2300
SACHAMENTO HIVER AT VEHONA	55055	1929-	USGS-DWR	3- 1-40	12.6	2,240	1 = 1 7 = 7 A	10.83	1930
SACRAMENTO WEIN SPILL TO YOLO PYPASS NEAR SACRAMENTO		1926=	USGS=NwP	3-26-28 12-23-55	10.0	3,340(86)			ND FLOW
NURTH FORK AMERICAN RIVER AT NUMBER FORK DAM	885	1941-	USGS	12-23-64	3.6	1,850	1 -5-78	1.50	272
RUMICUM WIVER NEAW FORESIMILE	A15	1458-	usas	12-23-64	16.9(41)		1-17-78	3.20	A5
MIDDLE FURK AMENICAN RIVER NEAR FORESTRILL	1 3 5 7	1958-	USGS	12-23-64	(14)0.15	A,780(I)	1-16-78	3.53	244
MIDDLE FORK AMERICAN RIVER NEAR AUHURN	1590	1911-	USGS	12-23-64	18.4(41)	7,160(1)	3 -4-78	4.08	237
SOUTH FORK AMERICAN RIVER NEAR CAMINO	1276	1922-	11868	12-23-55	14)9,9	1.410	6-15-78	1.91	2.4
SOUTH FURK AMERICAN RIVER NEAR LUTUS	1743	1951=	USGS	12-23-55	6.5	2.030	1-17-78	3.02	0.75
AMERICAN MIVEM AT HAIM (AMS (MEHIME HILSUM DAM)		1904-55	บรคร	11-21-50	9.7(0)	2,030			
AMERICAN HIVEH AT FAIR DAKS (AFTER FOLSOM DAM)	4884	1955=	USGS	12-23-64	h.6	3,260	5 =5=7A	3.02	245
SACRAMENTO HIVEN AT SACRAMENTO	60942	1879-	USGS=DAF	11-21-50	9,2(0)	2,940	3 -9-78	0.00	2250
SACHAMENIO HIVEH AT HALMIT GHOVE		1929-	DAR	12-25-64	3.7		3 -9-78	3.01	0.0
ADUME THEFH NEAR MELSEYVILLE	15	1954-	USGS	12-22-64	2.A	42	1-14-78	2.34	26
MELSEY CHEEK	99	1946-	uses	12-21-55	3.4	249	1-16-78	3.44	136
WEAR TURER TORE	1367	1944-	USGS	2-24-58	2.9	556	4 -6-78	2.33	125

1 STREAM AND STATION	APEA IN	O.E.	SOURCE	•	VIDUS MAXIM OF RECORD		•	1977-19 WATER Y	FAR
1	SO MILES	RECORD	RECORD	. DATE .	. STAGE . . IN FEET .	DISCHARGE IN CFS	. DATE .	STAGE IN FEET	. DISCHARGE
					AREA (CONTI				•••••••
SACRA**F	VTO RIVER VUED)	HASIN							
FEATHER HIVER AT YUMA CITY	3974	1943-	USGS=DWP	12=23=64 12=24=55	76.4 82.4	172,000	3 -7-78	49.89	
MORTH YUBA HIVER BELUM GODOYEARS HAR	250	1930=	USGS	2- 1-63	23.8(A)	40,000	1=16=78	9.00	4550
NORTH YURA KIVEK HELUW NEW HULLARDS HAR DAM NEAR SAN JUAN	490	1940=	USGS	1-22-70	35.3 40.5(C)	56,200 91,600(M)	12-15-77	5.97	030
SOUTH YUMA HIVEH NEAM CISCO	52	1942-	USGS	1-31-63	20.6(4)	18,400	5-14-78	7.33	2100
SOUTH YUMA HIVEF AT JONES HAR NEAR GRASS VALLEY	30 A	1940-48	USGS	12=22=64	25.0(4)	53,600	1 -9-78	12.70	8250
ANDFERMICH LOW	1108	1941-	USGS	12-22-64	564,1(0)	171,000(K)	1 -5-78	15.67	16200
DEER CHEEK NEAM SMAMIVILLE	A5	1935-	USG8	10-13-62	13.8	11,600	1 =5=78	8.58	3750
YLIAA RIVEH NEAH MARYSVILLE	1339	1940=	USGS	12=22=64	90.2	180,000	1-16-78	69.83	18700
MEAR WHEATLAND	292	1928=	USGS	12-22-55 11-21-50	19.3(C) 20.8(C)	33,000	1-17-78	13.11	9000
FFATHER RIVER AT NICHLAUS	5920	1943-	USGS-0WP	12=23=55	51.6	357,000	1 -6-78	33.40	26200
FREMONT WELK (MEST END) SPILL TO MOLD HYPASS		1934-	[546]	12=23=55	39.7	294,000(8)	1-18-78	36.43	81400
SACRAMENTO FIVER AT VERONA	21257	1929=	USGS-AWR	3- 1-40	41.2	79,200	1-17-78	35.52	68300
SALHAMENTO WEIH SPILL TO YOUR HYPASS MEAR SACHAMENTO		1926=	USGS=DWR	3=26=28 12=23=95	32.A 33.0	118,000(8E			NO FLOW
NORTH FORK AMERICAN RIVER AT NUMBER FURK DAM	342	1941-	USGS	12-23-64	11.9	65,400	1 -5-78	4.93	9650
RUBICUN HIVER MEAR FURESTRILL	315	1958-	USGS	12=23=64	55.4(41)		1-17-78	10.50	3000
MINDLE FORESTHILL	524	195#=	USGS	12=23=64	69.0(41)	310,000(1)	1-16-78	11.59	8650
MIDDLE FIFK AMERICAN RIVER NEAR AUBLING	614	1911-	USGS	12=23-64	60.4(AI)	253,000(1)	3 -4-78	13.38	8400
SOUTH FORK AMERICAN RIVER NEAR CAMINO	493	1922-	USSS	12-23-55	32.6(A)	49,800	6-15-78	6.26	080
SCHITH FORK AMERICAN PIVER NEAR LUTUS	673	1951-	USGS	12-23-55	21.4	71,800	1-17-78	9.91	8150
AMERICAN WIVEN AT FAIR DAKS (HEFDRE FOLSUS DAM)	1848	1904=55	USGS	11-21-50	31.9(0)	180,000			
AMERICAN HIVER AT FAIR HAKS (AFTER FOESHM DA")	1848	1955-	uses	12-23-64	21.6	115,000	5 =S=7A	9.97	8650
SACRAMENTO PIVER AT SACRAMENTO	23530	1879=	USGS-OWR NOAA	11-21-50	30,1(0)	104,000	3 -9-78	0.00	79300
SACHAMENTO RIVER		1929=	Or R	12-25-64	12.2		3 -9-78	9.89	
ADDRE CREEK NEAR KELSEYVILLE	6	1954-	USGS	12-22-64	9.1	1,500	1-14-78	7.69	950
MELSKY CHEEK NEAP MELSEYVILLE	3.7	1946-	USGS	12=21=55	12,8	8,800	1=16=78	11.29	4800
CACHE CHEEK LAFE	528	1944-	USGS	2-24-58	9.4	8,000	4 -6-78	7.63	4400

I I I STMFAM AND STATION	DRAIMAGE			PRE	OF RECORD				
	SO KM	. PECORO	RECORD	DATE	STAGE . IN METRES .	DISCHARGE .	DATE .	STAGE .	DISCHARGE IN 4 /S
			CENT	RAŁ VALLEY	AREA (CONTIN	(UED)			
SACRAME! (CUNTIF	NEVIO PIVER	PASTN							
NORTH FORE CACHE CHEEK	510	1930-	usas	12-11-37	4.3(A)	574	1-16-78	2.39	99
ACHE CHIEK AT HUMSEY	2473	1960-	US(,S=D+R	1= 5=65	6,5(AC)	1,670	1-16-78	6.21	424
CACHE CHEEK AT YHLH	2949	1903-	USGS	2=25-58 3=10-04	26.0 26.9(P)	1,170	1-16-78	22.17	532
YULD HYPASS NEAH ADUULAND		1939-	USGS+DWR	2- A-42	9.8	7,700	1-18-78	A.43	2420
PUTAH CREEK NEAH WINTERS	1486	1430-	USGS=DWR	2-27-40	9.3	2,290	1-14-78	2.85	31
YOLO HYPASS NEAH LISHUM		1914-	DWR	12-25-64	7.5	9,910(E)	1-19-78	5.77	0.0
SACRAMENTO WIVER AT WIO VISTA		1906-	DWR	12-26-55	3+1	(D)	6-30-78	2.40	0.0
SAN JOAN	JUIN PIVE	R HASIN							
RILLOW CHEEK AT MOUTH NEAH AUHERRY	337	1952•	uses	12-23-55	A.7(A)	444	2 -9-78	4.55	124
SAN JHAGHIA MIVER MELOW KEMCHUEM POMEMHOUSE NEAM PRATHEM	5A35	1942-	usgs	12-23-55	15.5(4)	2,610	6-10-78	7.73	438
SAN JUADUIN FIVER	4340	1907-	USGS	12-11-37	7.3(CM) 3.6	2,190(M) 351	4-26-78	3.00	216
SAN JURUUN MIVER NEAH MEUUNAM HABN	11163	1939-	USPR=NWR	6- 1-52 6-20-41	4.2(C)	250 332(M)	5 -6-7A	4.22	1 4 3
FRESHI HIVER NESH KMINES	344	1911-13	USGS	12-23-55	3.5	376	1-16-78	2.19	108
FRESNU HIVEH Near Daue Tou	668	1941-	USGS	12=23=55	3.A	495	2-14-7A	2.52	80
CHONCHILEA HIVEN HELOW HAYFOR CHEEK DEAR HAYMON	657	1972=	uses	2=11=73	3.0	314	4-25+78	2.65	91
EASTSIDE MYMASS REAM EL NION		1964-	Dww	2=25=69	5.4	614	4-26-78	5.08	aar
MERCEU HIVEN AL PUHDAGO HIDGE HEAR YOSEMITE	A31	1916-	USGS	12-23-55	6.6(4)	662	6 -9-7A	3.11	182
MERCEO RIVER DEAR STEVINSON	3297	1940-	USGS	12- 5-50	22.5	0,385	STATION D	1 SCONT I NU	E D
SAN JHANLIN HIVEH	24657	1912-	USGS=DHR	3-26-69	20.1(4)	982(L)	4-10-78	19.66	437
URESTIMBA CHEFM NEAR DEVISAN	347	1932=	เลลยบ	4- 2-5A	2.0(0)	885	1-17-7A	2.45	123
SOUTH FORK THOLUMME RIVER NEAR TORLAND HECHEATIUM CAMP	225	1923=	uses	12=23=55	3,3(4)	336	2 -9-78	2.42	65
HAMIJIII ATUTU PLATI MARAND PERMENTINI CAMP	191	1916=	USGS	12-23-55	3,6(A)	139	2 -9+7A	1.86	28
THDEHMNE PIVER AT MUDESTE	4879	1040-	USGS=0#R	12= 9=50	21.1	1,610	5 -7-7A	14,82	166

D.	2474465		SOUPCE	. PRE	VIOUS MAXIM			1977-1978	
STREAM AND STATION . AF	REA IN	. OF	. OF . RECORD		STAGE . 1N FEET .		DATE .	STAGE . IN FEET .	
,		• • • • • • • • • • • • • • • • • • • •	CENT		AREA (CONTI		••••••		
SACRAMEN (CONTINE		R RASIN							
ORTH FORK CACHE CREEK NEAR LOWER LAKE	197	1930-	USGS	12-11-37	14.0(A)	20,300	1-16-78	7.85	3500
ACHE CREEK AT HUMSEY	955	1960-	USGS-DWR	1- 5-65	21.4(AC)	59,000	1-16-78	20.37	15000
ACHE CREEK AT YOLO	1139	1903=	USGS	2-25-58 3-10-04	85.4 88.4(P)	41,400	1-16-78	72.73	18800
OLO BYPASS NEAR WOODLAND		1939•	USGS=0W9	2- 8-42	32.0	272,000	1-18-78	27.67	85600
UTAH CREEK NEAR WINTERS	574	1930-	USGS-DWR	2-27-40	30.5	81,000	1-14-78	9.34	1150
OLO BYPASS NEAR LISHON		1914-	OWR	12-25-64	24.7	350,000(E)	1-19-78	18.94	
ACRAMENTU HIVER AT HIO VISTA		1906-	DWR	12-26-55	10.2	(0)	6-30-78	7.86	
SAN JOAG	nin ein	ER BASIN							
ILLOW CHEEK AT MOUTH NEAR AUHERRY	130	1952*	USGS	12=23=55	28.5(A)	15,700	2 -9-78	14.92	4400
AN JOAQUIN MIVER HELOW KERCHOFF POWERHOUSE NEAR PRATHER	1481	1942-	USGS-	12-23-55	51.0(A)	92,200	6=10=78	25,35	15500
AN JOAQUIN PIVER BELUW FRIANT	1676	1907-	USGS	12-11-37	23.8(CM) 11.7	77,200(M) 12,400	4-26-78	9.83	7650
AN JUADUIN HIVER NEAR MENDUTA	4310	1939-	USBR+DwR	6= 1=52 6=20=41	13.8(C)	8,840 11,740(M)	5 =6=78	13.86	5050
RESNO PIVEH NEAH KNOWLES	133	1911=13 1915=	USGS	12-23-55	11.5	13,300	1-16-78	7.18	3800
RESNO RIVER NEAR DAULTIN	258	1941-	USGS	12-23-55	12.6	17,500	2-14-78	8.27	2850
HOWCHILLA PIVER HELDW RAYNOR CREEK NEAR RAYMOND	254	1972-	USGS	2=11=73	9.9	11,100	4-25-78	8.69	3250
ASTSIDE HYPASS NEAP EL NICO		1964-	OwR	2-25-69	17.6	21,700	4-26-78	16.68	15800
ERCED RIVER AT POHONO BRIDGE NEAR YOSEMITE	321	1916-	USGS	12-23-55	21.5(A)	23,400	6 -9-78	10.19	6450
EMCED RIVEW NEAR STEVINSON	1273	1940-	USGS	12- 5-50	73.A	13,600	STATION	OISCONTINUE	
NEAR NEWWAY	9520	1912-	USGS=0WR	2=26=69	65.9(4)	34,700(L)	4-10-78	64.49	15400
RESILMHA CHFEK NEAR NEWFAN	134	1932=	USGS	4- 2-58	6.6(C)	10,200	1-17-78	8.03	4350
OUTM FORK TOULDMAF RIVER NEAR CLAKLAND RECREATION CAMP	A 7	1923-	USGS	12-23-55	10.9(A)	11,900	2 -9-78	7.90	2300
TODLE THOLLIMNE RIVER AT CAKLAND RECKEATION CAMP	74	1916=	USGS	12-23-55	11.8(A)	4,920	2 -9-78	6.09	1000
HOLUMME PIVER	1884	1940=	USGS=0wP	12- 9-50	69.2	57,000	5 -7-78	48.63	5911

I I	ORAINAGE	. PER10D	ERIOD . SDURCE		EVIOUS MAXIM		WATER YEAR			
1 STREAM AND STATION I	AREA IN	. DF . RECORD	. DF . RECORD :	DATE .	STAGE . IN METRES .	DISCHARGE .	DATE .	STAGE IN METRES	. DISCHARGE	
			CEN71		AREA (CONTTI				••••••	
SAN JO	AQUIN RIVE INUED)	R BASIN								
SDUTM FORK STANISLAUS RIVER NEAR LONG BARN	173	1937=	USGS	11-21-50	8.5	138	7 -3-78	1.76	36	
STANISLAUS HIVER AT DRANGE HLUSSUM BRIDGE		1928-39 1940-	DWR	12-23-55	9.7	1,760	5-23-78	3.47	152	
STANISLAUS RIVER AT RIPDN	2784	1940-	USGS-0wR	12-24-55	19.3 19.6(A)	1,770	5 -8-78	15.51	113	
SAN JOAQUIN RIVER NEAR VEHNALIS	3506R	1922+	USGS-DWR	12- 9-50	10.0(C) 10.5	2,240 1,490	5 -4-78	8.57	747	
DUCK CREEK NEAR STUCKTON		1950-	DMR	1-16-73	5.0	55	3 -4-78	1.81	17	
SOUTH FURK CALAVERAS RIV NEAR SAN ANDREAS	305	1950=	USGS	12-23-55	3.1	498	1-14-78	2.51	123	
MORMUN SLOUGH AT BELLOIA		1948-	DWR	4- 2-58	6.3	436(E)	2 -7-78	2.74	49	
STOCKION DIVERTING CANAL AT STUCKION		1944-	DwR	4- 4-58	5.2(E)	355(E)	1=15=78	3,33	107	
CALAVENAS HIVEH NEAR STOCKTON		1958-	DWR	1- 6-65	3.8	21(E)	2 -8-78	1.92	6.4	
NEAR LOCKEFORD	124	1930-	USGS	4- 3-58	4.6	82	1+14-7A	2,69	37	
CDLE CHEEK NEAR SALT SPRINGS DAM	52	1927-42	USGS	12-23-64	3.1	173	5-14-78	1.18	35	
SOUTH FUNK MOKELUMNE RIV NEAR WEST POINT	/FR 194	1933-	USGS	12-23-55	4.5(AC)	195	1-14-78	1.93	34	
MEAR MORELUMNE HILL	140 R	1901-	USGS	12- 3-50	5.6	954	6 -8-7R	3,95	170	
MOKELUMNE RIVER	1711	1924-	USGS	11-22-50	9.0	764	5 =3=7R	3,69	38	
NR THORNTON (RENSON FERS	71) 5296	1911-	DMR=NDAA	12-24-55	5.5(C)	- ~(D)	1-16-78	3.06	0.0	
NEAR GALI	852	1926-33	USGS-DWR	4- 3-58	4.7	679	3 -6-78	4.21	142	
NORTH FORK COSUMNES RIVE NEAR EL DURADU	- H 530	1948-	USGS	12-23-55	4.5	447	1-17-78	5.44	P-1	
SOUTH FORK COSUMMES RIVE MEAN PIVEN PINES	165	1957-	USGS	2- 1-63	3,3	156	1-15-78	1.70	53	
CUSUMNES PIVER AT MICHIGAN HAR	1388	1907-	USGS-DKR	12-23-55	4.5 5.0(A)	1,190	3 =4=7A	2.40	233	
AT MCCUNNELL	1875	1941-	U968	12-23-55	14.1	1,530	1-15-76	13.43	291	
	LAKE RAST	I N								
NEAR SPRINGVILLE	639	1957=	USGS	12= 6=66	6.0(AC)	1,400	2 -9-78	2.90	555	
TOLE MINER SUCCESS DAM	1017	1953-	usgs	12-23-55 11-19-50	6.6(C) 7.9(AC)	764 906(M)	3-16-78	2.03	55	
MAMEAM MIVEN AT THMEE MIVERS	1082	1958•	USGS	12- 5-66 12- 5-66	5.1 5.8(A)	2,070	3 -4-78	2.86	267	
RINGS RIVER RELUW NORTH FURK	3475 V15TA LAKE	1951=	USGS	12=23=55	7.0	2,410	9 -5-78	3.73	615	
(, 5, 1, 2,	10.2 6-00									

5.9(A) 2,100 6 -9-78 3.10

550

1905=12 2612 1953= USGS

FERN RIVER AT KERNVILLE

PEAK FLOWS AND STAGES (CONTINUED)

I .				PRE	VIDUS MAXIMU			1977-19	78 1
1 . 0	HAINAGE PEA IN	PERIOO OF	. SOURCE		OF RECORD			WATER Y	EAR 1
1 SINEAM AND SIBILON . A	WILES	RECORD	RECORD	. 041E .	STAGE .	DISCHARGE .	DATE .	STAGE	. DISCHARGE I
1 .					IN FEET .			IN FEET	
			CENTI	RAL VALLEY	AREA (CONTIN	(UED)			
DAUL MAP		R BASIN							
SDUTH FORK STANISLAUS RIVER NEAR LONG HARN	67	1937-	USGS	11-21-50	9.3	4,900	7 -3-78	5.76	1300
STANISLAUS HIVER AT DRANGE HLOSSOM BRIDGE		1928-39 1940-	0#R	12-23-55	31.8	62,000	5-23-78	11.39	5400
STANISLAUS HIVEH AT HIPON	1075	1940-	USGS-DWR	12-24-55 2+12-38	63.3 64.4(A)	62,500	5 -8-78	50.88	4000
SAN JOANUIN HIVER NEAH VERNALIS	13540	1922-	USGS•D#R	12- 9-50 1-27-69	32.8(C) 34.6	79,000 52,600	5 -4-78	28.12	26400
NEAR STUCKTUN		1950-	OwR	1-16-73	6.5	780	3 -4-78	5.94	600
SOUTH FURK CALAVERAS RIVER NEAR SAN ANDREAS	118	1950-	USGS	12-23-55	10.3	17,600	1-14-78	8,22	4350
MORMON SLOUGP AT BELLUTA		1948-	DwR	4- 2-58	20.7	15,400(E)	2 -7-78	8,98	3500
STOCKTON DIVERTING CANAL AT STOCKTON		1944-	DWB	4- 4-58	17.1(E)	11,400(E)	1-15-78	10.94	3800
CALAVERAS HIVER NEAR STOCKTON		1958=	DWR	1- 6-65	12.6	760(E)	2 -8-78	6.29	250
BEAR CREEK NEAR LOCKEFURD	48	1930-	USGS	4- 3-58	15.1	2,930	1-14-78	8.81	1300
COLE CREEK NEAR SALT SPRINGS DAM	20	1927-42 1943-	USG\$	12-23-64	10.2	6,140	5-14-78	3.87	1250
SOUTH FURK MUKELUMNE RIVER NEAR WEST POINT	75	1933-	USGS	12-23-55	14.8(AE)	6,920	1-14-78	6.34	1200
MUKELUMNE RIVEH NEAR MUKELUMNE HILL	544	1901-	USG\$	12- 3-50	18.5	33,700	6 -8-78	12.97	6000
MORFLUMNE RIVER AT WOODHRIGGE	661	1924-	USGS	11-22-50	29.6	27,000	5 =3=78	12.11	1350
MOKELUMNE HIVER NR THORNION (BENSON FERRY)	2045	1911-	OWR-MUAA	12-24-55	18.0(E)	(0)	1=16=78	10.05	
DRY CREEK NEAH GALT	329	1926-33	USGS=DWF	4- 3-5A	15.3	24.000	3 -6-78	13.81	5050
NORTH FORK CUSUMMES RIVER NEAR EL MORADO	205	1911-41	USGS	12-23-55	14.8	15,800	1-17-78	7.99	2900
SHUTH FORK COSUMNES RIVER NEAR HIVER PINES	64	1957•	USGS	2- 1-63	10.9	5,540	1-15-78	5.57	1900
COSUMPES MIVEN AT MICHICAN HAP	536	1907=	USGS-DWF	2 12-23-55 307	14,6 16,3(A)	42,000	3 =4=7A	7.88	8250
COSIMANES PIVER	724	1941-	USGS	12-23-55	46.3	54,000	1-15-78	44.07	10300
JIII ARE	LAFE HAS	IN							
THE RIVER SPHINGVILLE	247	1957=	USGS	12- 6-66	19.7(AC)	49,600	2 -9-78	9.52	7850

12-23-55

12= 5=66 12= 5=66

12-23-55

21.7(C) 26.0(AC)

> 16.7 19.0(A)

19.3(A)

23.1

27,000 32,000(M)

73,000

85,200

74,000

3-16-78

3 -4-78

9 -5-78

6 -9-78

6,67

9.37

12.24

10.18

800

9450

21700

7800

TOLF RIVER BELOW SUCCESS DAM

KAWFAH WIVEN AT IHREE MIVENS

KINGS HIVER BELUK NUFTH FURK

MEHN RIVER

303

418

1 542

1009

HILFRA VISTA LAKE PASIM

1953-

1958-

1951-

1905=12

USGS

USGS

USGS

HSGS

STHEAM AND STATION . API	PAINAGE . PEP1	PEPIND	PEPIND . SOURCE		OF RECORD		1977-1978 WATER YEAR			
	SQ KM	. MECORD	. RFCORD	. DATE .	STAGE .	DISCHARGE . IN M /S .	DATE .	IN METRES	. DISCHARGE . IN 4 /S	
				HERN LAHONT						
HONEY L	AKE HASIN									
ILLON CHEEK NEAM SUSANVILLE	233	1950-	USGS	2- 1-63	1.7	23	1-16-78	1.35	9,3	
SUSAN MIVEM AT SUSANVILLE		1917=21 1950=	USGS	12-22-64	5.5	144	1-14-78	1.34	20	
LAKES PYRAMIC	AND WINH	EMUCCA								
117LL THUCKEE RIVER AMON BUCA RESERVUIM NEAR BUCA		1903-10 1939-	USGS	2- 1-63	2.7	376	5-16-78	0.98	49	
NUCKEE HIVER AT FAMAD	2413	1499-	U868	11-21-50	4.4(A)	495	5-21-78	1.89	94	
CARSON	RIVER RAS	1 M								
AST FORK CARSON RIVER RELUW MARKLEEVILLE CREEK	714	1960-	USGS	1-31-63	3.1	427	5-15-78	1.30	66	
VEST FORK CARSUN RIVER AT WOODFORDS		1900-07	USGS	2- 1-63	2.7	138	5-14-78	1.14	25	
WALKER	LAKE PASI	N.								
NEST VALKEM RIVER HELUA LITTLE WALKER RIVER NEAM CULEVILLE	466	1938-	USGS	11-20-50	2.5	176	6-14-78	1.50	56	
MEAN HRIDGEPUHI		1911-14	USGS	6-19-63	1.4	39	5 -3-78	0.84	1 4	
			sour	THERM LAHONT	AN AREA					
MUJWAF	FIVER RAS	I M								
ANJAVE FIVER AT LUWER NARHUAS MEAR VICTORVILLE		1899-06 1930-	USGS	3- 2-3R	7.2	2,000	2-10-78	2.58	396	
MOJAVE FIVER	3341	1930-	USGS	3- 3-38	2.6	1,820	2-10-78	1.45	291	
#3V1 3VALO" AI)F14 TA		1952-32	USGS	1-26-69	3.2	509	3 -5-78	0.00	396(E	

1 1 . DF	RAINAGE . PERIUD	UD . SOURCE		OF RECORD		. 1977-1978 WATER YEAR			
I STREAM AND STATION . A)	MILES	. RECORD	. RECORD	. DATE .	STAGE . IN FEET .	DISCHARGE .	, DATE .	STAGE . IN FEET .	DISCHARGE I IN CFS I
				THERN LAHONT					
HONEY 1 A	KE BASI	N							
WILLOW CREEK NEAR SUSANVILLE	90	1950-	USGS	2= 1=63	5.6	820	1-16-78	4.43	350
SUSAN RIVER AT SUSANVILLE		1917+21 1950+	USGS	12-22-64	7.3	5,100	1-14-78	4.41	750
PYHAMIO AND WINNEMUCCA LAKES BASIN									
LITTLE TRUCKEE RIVER AROVE BOCA RESERVOIR NEAR BOCA			USGS	2= 1=63	9.0	13,300	5=16=78	3,21	1750
TRUCKEE HIVEH	932	1899=	USGS	11-21-50	14.5(A)	17,500	5-21-78	6.19	3350
CARSON R	IVER HA	SIN							
EAST FORK CARSON RIVER BELOW MARKLEEVILLE CREEK	276	1960-	USGS	1=31=63	10.2	15,100	5=15=78	4.27	2350
WEST FORK CARSON RIVER AT WOODFURDS		1900-07	USGS	2= 1=63	9.0	4,890	5-14-78	3.75	900
MALKER L	AKE BAS	IN							
WEST WALKER RIVER BELUW LITTLE WALKER RIVER NEAR COLEVILLE	180	1938+	USGS	11-20-50	A.1	6,220	6=14=78	4.92	5000
EAST WALKER RIVER NEAM BRIDGEPORT		1911-14	USGS	6=19=63	4.6	1,390	5 -3-78	2.76	500
			8011	THERN LAHON1	AN AREA				
MOJAVE R	IVER RA	SIM							
MOJAVE RIVER AT LOWER NARRIWS NEAR VICTORVILLE			usgs	3- 2-38	23.7	70,600	2-10-78	8.48	14000
MUJAVE RIVER AT HAMSTOW	1290	1930=	USGS	3= 3=38	8.6	64,300	2-10-78	4.75	10300
MUJAVE RIVER AT AFTON	2120	1929=32 1952=	USGS	1=26=69	10.4	18,000	3 +5=78	0.00	14000(E)

LEGEND

- USGS United States Geological Survey
- USBR United States Bureau of Reclamation
- NOAA National Weather Service (National Oceanic and Atmospheric Administration)
- USCE United States Corps of Engineers
- Department of Water Resources DWR
- PG&E Pacific Gas and Electric Company
 - A From flood marks
 - B Discharge over weir or spillway
 - C Site or datum then is use
 - D Discharge not determined, affected by backwater or tide
 - Ε Estimated
 - F From DWR telemetering log
 - G Preliminary
 - H Includes flow through power plant
 - I Due to failure of partially completed dam
 - Gage height revised
 - K Flow through power plant not included
 - L Discharge at latitude of gaging station site
 - M Prior to construction of upstream dam
 - Includes flow through fish hatchery but not upstream N diversion to Thermalito Afterbay
 - P Observed
 - Q Estimated peak inflow to partially completed Oroville Reservoir
 - R Regulated stage and flow
 - Revised to current datum
 - T Datum of gage is 0=0 USED
 - U Crest stage partial recorder
 - N/A Not available at report time
 - Peak of record established current year





, * Mov 51 .

1131-31053

4 - L1 1 . Y

JUN 30 1984

FILES

3 0 1984

日本 5. 1 11 11 11 11



State of California—Resources Agency
Department of Water Resources
P.O. Box 388
Sacramento
95802





JUN 7 1980

APR 6 '84

CALIFORNIA, UNIVERSITY OF DAVIS CAMPUS LIBRARY, GOVERNMENT DOCUMENTS OFP DAVIS, CA 95616

بالاستادات





